

A Connection between Pre-mRNA Splicing and the Cell Cycle in Fission Yeast: *cdc28*⁺ Is Allelic with *prp8*⁺ and Encodes an RNA-dependent ATPase/Helicase

Karen Lundgren,^{*†} Susan Allan,^{*} Seiichi Urushiyama,[‡] Tokio Tani,[‡] Yasumi Ohshima,[‡] David Friendewey,[§] and David Beach^{*||}

^{*}Howard Hughes Medical Institute, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 11724; [‡]Department of Biology, Faculty of Science, Kyushu University, Hakozaki, Higashi-ku, Fukuoka 812-81, Japan; and [§]Department of Microbiology, New York University School of Medicine, New York, New York 10016

Submitted March 12, 1996; Accepted May 8, 1996
Monitoring Editor: Mitsuhiro Yanagida

The fission-yeast gene *cdc28*⁺ was originally identified in a screen for temperature-sensitive mutants that exhibit a cell-division cycle arrest and was found to be required for mitosis. We undertook a study of this gene to understand more fully the general requirements for entry into mitosis. Cells carrying the conditional lethal *cdc28-P8* mutation divide once and arrest in G2 after being shifted to the restrictive temperature. We cloned the *cdc28*⁺ gene by complementation of the temperature-sensitive growth arrest in *cdc28-P8*. DNA sequence analysis indicated that *cdc28*⁺ encodes a member of the DEAH-box family of putative RNA-dependent ATPases or helicases. The Cdc28 protein is most similar to the Prp2, Prp16, and Prp22 proteins from budding yeast, which are required for the splicing of mRNA precursors. Consistent with this similarity, the *cdc28-P8* mutant accumulates unspliced precursors at the restrictive temperature. Independently, we isolated a temperature-sensitive pre-mRNA splicing mutant *prp8-1* that exhibits a cell-cycle phenotype identical to that of *cdc28-P8*. We have shown that *cdc28* and *prp8* are allelic. These results suggest a connection between pre-mRNA splicing and progression through the cell cycle.

INTRODUCTION

Intensive genetic and biochemical studies of the cell-division cycle have converged to give a relatively cohesive picture of the major eukaryotic cell-cycle regulators.¹ In yeasts, control of the cell cycle is regulated primarily by the activity of the protein kinase p34^{cdc2} (Draetta, 1990; Nurse, 1990). This activity is, in turn, regulated by physical association with a family of proteins known as cyclins (Draetta, 1990; Pines and Hunter, 1991) and by phosphorylation and dephos-

phorylation at specific and critical points in the cell cycle (Draetta *et al.*, 1988; Morla *et al.*, 1989; Ducommun *et al.*, 1991; Gould *et al.*, 1991).

Many cell-cycle regulators have been identified in genetic screens for cell-division cycle (*cdc*) mutants in the budding yeast *Saccharomyces cerevisiae* (Hartwell, 1978) and the fission yeast *Schizosaccharomyces pombe* (Nurse, 1975). These mutations define genes whose products are required for progression through the cell cycle. In many cases these proteins have been shown to have conserved functions in all eukaryotes. The *S. pombe cdc2*⁺ gene was identified in such a screen and shown to be required at two points in the cell cycle: in G1, before start, and in G2 (Nurse and Bissett, 1981). Several genes have been identified that regulate p34^{cdc2} activity. These in-

[†] Present address: Glaxo Wellcome, 5 Moore Drive, Research Triangle Park, NC 27709.

^{||} Corresponding author.

¹ The GenBank accession number for the sequence reported in this manuscript is U48733.

clude *wee1⁺* and *mik1⁺*, a pair of tyrosine kinases, and *cdc25⁺* and *pyp3⁺*, a pair of tyrosine phosphatases, which function antagonistically to regulate the phosphorylation state of p34^{cdc2} (Russell and Nurse, 1986, 1987; Lundgren *et al.*, 1991; Millar *et al.*, 1992). In addition, four cyclins —*cdc13⁺*, *puc1⁺*, *cig1⁺*, and *cig2⁺* —that function as regulatory subunits of the p34^{cdc2} protein kinase have been identified (Booher and Beach, 1987, 1988; Bueno *et al.*, 1991; Bueno and Russell, 1993; Forsburg and Nurse, 1991; Connolly and Beach, 1994).

Other *cdc* genes in fission yeast define functions that, although not directly related to regulation of p34^{cdc2}, must be executed for the cell cycle to progress. Examples include *cdc17⁺*, the gene for DNA ligase (Nasmyth, 1977), *cdc22⁺* and *suc22⁺*, which encode the large and small subunits of ribonucleotide reductase (Fernandez Sarabia *et al.*, 1993), and *cdc10⁺* and *sct1⁺*, which encode components of a transcription complex required for commitment to the cell-division cycle at start (Aves *et al.*, 1985; Lowndes *et al.*, 1992; Caligiuri and Beach, 1993). Thus, mutations in genes that regulate p34^{cdc2} as well as in genes that participate in general cellular functions, such as DNA synthesis and transcription, can elicit *cdc* phenotypes.

Mutations in genes involved in RNA metabolism can also affect cell-cycle progression. A prime example is the *pim1⁺* gene (Matsumoto and Beach, 1991), whose mammalian homologue RCC1 encodes a protein that prevents premature initiation of mitosis but also participates in nuclear RNA processes (Dasso, 1993; Rush *et al.*, 1996). Mutations in the *Saccharomyces cerevisiae* homologue of RCC1 cause global defects in nuclear structure, RNA synthesis, and transport (Aebi *et al.*, 1990). A specific relationship between pre-mRNA splicing and cell-cycle progression is illustrated by the *dbf* mutants of *S. cerevisiae*. The *DBF3* and *DBF5* loci were identified in a screen for DNA synthesis mutants blocked in S phase, but they were found to be allelic with *PRP8* and *PRP3*, two genes required for pre-mRNA splicing (Johnston and Thomas, 1982; Shea *et al.*, 1994). These and other emerging examples point to a connection between RNA metabolism and regulation of the cell-division cycle.

The *S. pombe cdc28⁺* gene was initially identified in a screen for *cdc* mutants and was found to be required for entry into mitosis (Nasmyth and Nurse, 1981). In an effort to define more fully the requirements for cell-cycle progression, the present study describes the isolation and characterization of the *cdc28⁺* gene. The unexpected results of this investigation support a role for pre-mRNA splicing in the regulation of the cell cycle.

MATERIALS AND METHODS

Yeast Strains and Growth Conditions

All *S. pombe* strains used in this study were derived from those originally described by Leupold (1970) and are listed in Table 1. *S. pombe* was grown in standard YEA, YE, and phorbol 12-myristate 13-acetate (PMA) media (Beach *et al.*, 1985) containing additional amino acids at 75 μ g/ml. In some cases phloxin B was added to solid medium at 20 μ g/ml as an indicator of cell viability. So that the growth rate and viability of the *cdc28-P8* strain could be determined, mutant and wild-type cells were grown at 25°C in rich media (YEA) to mid-log phase and then shifted to the restrictive temperature (37°C). Cells were counted and replated on YEA at 25°C at the start of the shift to 37°C and at hourly intervals thereafter. Relative viability was calculated as the number of colonies at a given time point compared with the number at the start of the shift to 37°C. For fluorescence-activated cell-sorting (FACS) analysis, cells were grown in PMA containing 0.6 mM NH₄Cl for 48 h to arrest the cells in G1 and then inoculated into YEA at 37°C. Cells were removed at 2-h intervals and prepared as previously described (Costello *et al.*, 1986). For 4,6-diamidino-2-phenylindole (DAPI) staining, 10⁷ cells grown to mid-log phase in liquid culture were washed in 5 ml of cold, deionized water. The cells were fixed in 1 ml of 2.5% glutaraldehyde for 10 min on ice and then washed twice in 1 ml of cold water. After resuspension in 25 μ l of cold water, a mixture of 5 μ l of cells and 5 μ l of 75 μ g/ml DAPI was prepared. A 2- μ l sample was placed on a slide and observed with an Olympus PM-10AK fluorescence microscope with a DAPO 100 UV objective.

Cloning of *cdc28⁺*

Standard molecular-cloning procedures were used (Sambrook *et al.*, 1989). We transformed (Ito *et al.*, 1983) SP968 with a genomic library (Caligiuri and Beach, 1993) constructed in pWH5 (Wright *et al.*, 1986). Transformants of *leu⁺* (3.2 × 10⁴) were obtained at 25°C. The transformants were replica plated to YEA containing phloxin B and grown at 37°C to select for temperature-resistant (*ts⁺*) clones. Eleven *ts⁺* transformants were tested for co-stability of the leucine prototrophy (*leu⁺*) and the ability to grow at 37°C. Of the plasmids recovered from the *ts⁺* transformants (Beach *et al.*, 1982), four were able to complement the growth defect when reintroduced into SP968. Restriction analysis of these plasmids revealed that all four carried the same genomic fragment. The 7.5-kb insert from plasmid pcdc28-1 was subcloned, and the complementing activity was retained on a 4.8-kb *Bam*HI-*Bgl*II fragment that contained part of the pWH5 vector. Overlapping deletions of this fragment were constructed (Henikoff, 1987), and single-stranded DNA was prepared (Vieira and Messing, 1987). The sequence of both strands was determined with a 373A-automated sequencer (Applied Biosystems, Foster City, CA). An open reading frame (ORF) of ~3 kb was found, and the predicted amino acid sequence was compared with other proteins in the database by using the BLAST program (Altschul *et al.*, 1990). The complementing activity was shown to carry the *cdc28⁺* gene and not a multicopy suppressor by integration mapping. The *ars* sequence was excised from pcdc28-1 by digestion with *Eco*RI. A *leu⁻* strain was transformed to leucine prototrophy with

Table 1. *S. pombe* strains

972	<i>h^{-S}</i>
SP793	<i>h^{-S} cdc28-P8</i>
SP968	<i>h^{+N} cdc28-P8 leu1-32</i>
JP23	<i>h^{-S} prp1-1 ade6-216</i>
DK633	<i>h^{-S} prp5-1 leu1-32</i>
SU50-5B	<i>h^{-S} prp2-2</i>
SU26-9B	<i>h^{-S} prp8-1</i>

the plasmid, and stable transformants, because of chromosomal integration, were isolated. One of these integrants was crossed to a *cdc28-P8 leu1-32* strain to determine whether integration had occurred at the *cdc28* locus. Twenty-one tetrads were dissected, and in each case the *leu* and *ts* markers segregated 2:2. All *leu*⁺ segregants grew at 37°C, and all *leu*⁻ segregants were *ts*⁻. We therefore conclude that the gene we isolated integrated at the *cdc28* locus and is the authentic *cdc28*⁺ gene. For genomic mapping the *Mlu* I fragment of the *cdc28*⁺ gene was labeled with ³²P by random priming and used to probe an *S. pombe* genomic cosmid library on filters (Mizukami *et al.*, 1993).

Isolation of *prp8-1*

The *S. pombe* pre-mRNA splicing mutant *prp8-1* was isolated from a bank of temperature-sensitive lethal strains made by ethyl methane sulfonate (EMS) mutagenesis. Initial screening of the bank for the mutants defective in pre-mRNA splicing was done by hybridization of Northern blots with probes complementary to U6 small nuclear RNA. The *S. pombe* gene for U6 RNA is highly expressed and has an intron similar in structure to nuclear pre-mRNA introns (Tani and Ohshima, 1989). Defects in pre-mRNA splicing in each candidate were then confirmed by Northern blot analysis with the use of a probe for the β -tubulin mRNA whose gene has five introns (Hiraoka *et al.*, 1984). Of 120 *ts*⁻ strains screened, we identified three mutants defective in pre-mRNA splicing. Complementation analysis with those mutants and previously isolated *prp* mutants (Potashkin *et al.*, 1989; D. Kim and D. Frendewey, unpublished results) showed that one of the mutants we isolated belongs to a new complementation group designated as *prp8*.

Genetic Linkage between *cdc28* and *prp8*

Strains carrying the *cdc28-P8* allele were crossed with strains carrying one of nine different *ts*⁻ pre-mRNA splicing alleles: *prp1-1*, *prp2-1* (Potashkin *et al.*, 1989); *prp3-2*, *prp4-2*, *prp5-1*, *prp6-1*, *prp7-1*, *prp9-1* (Kim and Frendewey, unpublished data), and *prp8-1* (S. Uemura and T. Tani, unpublished data). For complementation grouping, diploid zygotes were isolated on rich medium at 23°C and tested for viability at 37°C. For linkage analysis, zygotes produced by the crosses were sporulated on maltose extract plates at 23°C. Unmated cells were killed and spores released by treatment with Glusulase (DuPont, Wilmington, DE). Random spores (800–2000) were germinated on a rich medium at 23 and 37°C, and the percentage of viable spores that grew at 37°C was determined.

Northern Analysis

Cultures were grown in rich media at 25°C to mid-log phase. Aliquots of the cultures were removed (time 0), and the remainder was shifted to 37°C. Samples were collected at 2, 4, 6, and 8 h after the temperature shift. Cell pellets were washed in RNase-free water and frozen at -70°C. The cells were lysed by vortexing with glass beads in 1 ml of solution A (0.32 M sucrose, 20 mM Tris-HCl, pH 7.5, and 10 mM EDTA) containing 0.5 mg/ml heparin. After centrifugation to remove the glass beads, the supernatant was transferred to a new tube and brought to 8 ml with solution B (50 mM Tris-HCl pH 7.5, 100 mM NaCl, 5 mM EDTA, and 1% SDS). The lysate was extracted twice with phenol and once with phenol/chloroform and ethanol precipitated twice. For analysis of pre-U6 RNA splicing, 10 μ g of RNA from each time point was fractionated by electrophoresis on a 10% polyacrylamide/7 M urea gel. The RNA was transferred to Hybond (Amersham, Arlington Heights, IL) and fixed according to the manufacturer's instructions. The blot was hybridized with ³²P-labeled I-2 and E2-2 oligodeoxynucleotide probes against intron and second exon sequences of the U6 RNA precursor, as previously described (Potashkin *et al.*, 1989). For analysis of pre-mRNA splicing, 20 μ g of total RNA was electrophoresed on a 1% formaldehyde agarose gel (Sambrook *et al.*, 1989), blotted onto a Gene Screen

membrane (DuPont-NEN, Boston, MA) by the capillary method, and fixed by UV irradiation. The membrane was probed with a mixture of two ³²P-labeled oligonucleotide probes (TFII-IN1 and TFII-EX3) complementary to the first intron or the third exon of the *S. pombe* transcription factor IID (TFIID) pre-mRNA (Hoffmann *et al.*, 1990) in a solution containing 6 \times SSC, 50 mM sodium phosphate (pH 7.5), 2 mM EDTA, 0.1% SDS, 5 \times Denhardt's solution, and 0.2 mg/ml of preboiled salmon sperm DNA at 42°C. After hybridization, the membrane was washed three times in 6 \times SSC at room temperature and then once in 6 \times SSC and 0.1% SDS at 50°C for 5 min. Autoradiography was performed at -70°C with an intensifying screen for 2 d. Nucleotide sequences of the oligodeoxynucleotide probes for TFIID mRNA are as follows: TFII-IN1, 5'-GAAATCTCGTGACATGGTAG-3'; and TFII-EX3, 5'-GAGCTTGGAGTCATCCTCGG-3'.

RESULTS

The *cdc28-P8* Mutation Causes Cell-Cycle Arrest in G2

Growth analysis of the *cdc28-P8* mutant indicated that it doubled once and then arrested upon shift to the restrictive temperature of 37°C (Figure 1A), and the cells became inviable (Figure 1B). Fluorescence-activated cell sorter (FACS) analysis demonstrated that *cdc28-P8* mutant cells grown in rich media accumulate in G2 when shifted to 37°C. In addition, we found that when *cdc28-P8* cells previously arrested in G1 by starvation for nitrogen were released into rich media, they arrested with a 2C-DNA content (Figure 1C, panel d). Because the fission-yeast cell cycle has a relatively long G2, most of the wild-type cells were also found to be in G2 (Figure 1C, panel c). These results demonstrate that *cdc28-P8* is able to proceed through G1 and S phases before arresting in G2, confirming the original observation of Nasmyth and Nurse (1981) that the *cdc28*⁺ gene product executes its function in G2. To characterize the phenotype of the arrested cells, *cdc28-P8* was examined cytologically by staining with DAPI. Wild-type and mutant cells were grown at 25°C and then shifted to 37°C. Cells were collected at 0, 4, and 8 h after the shift to 37°C (Figure 1D). After 4 h at the restrictive temperature, the mutant cells (Figure 1D, panel b) were elongated and began to show cell-cycle arrest; after 8 h at 37°C, the cells were more elongated, and the nuclear material seemed to be condensed (Figure 1D, panel c). Wild-type cells at 37°C maintained the normal cellular and nuclear morphology (Figure 1D, panel d).

The *cdc28*⁺ Gene Encodes an RNA-dependent ATPase/Helicase

We cloned the *cdc28*⁺ gene by complementation of the temperature-sensitive (*ts*⁻) growth impairment of *cdc28-P8*. A partial *Hind*III-*S. pombe* gene bank (Cali giuri and Beach, 1993) on a replicating plasmid carrying the *S. cerevisiae* *LEU2* gene, which complements *leu1-32*, was introduced into SP968 (*cdc28-P8 leu1-32*). We selected *leu*⁺ transformants that grew at 37°C and




Figure 1. Phenotype of the *cdc28-P8* mutant. (A) Growth rates of *cdc28-P8* compared with the wild-type strain. Cells from cultures of exponentially growing *cdc28-P8* (SP793) and wild type (972) were counted at intervals after shifting to 37°C. (B) Cell viability. The relative viability of *cdc28-P8* and wild-type cells after shift to the restrictive temperature was determined by the ability to form colonies at 25°C. (C) FACS analysis. *cdc28-P8* and wild-type cells were grown in minimal media with limiting nitrogen at 25°C to starve the cells and arrest them in G1. The cells were then released into rich media at 37°C. Samples were removed for FACS analysis before and at 4 h after the temperature shift. Panels: (a) Wild type before the shift; (b) *cdc28-P8* before the shift; (c) wild type after 4 h at 37°C; and (d) *cdc28-P8* after 4 h at 37°C. (D) Fluorescence microscopy of DAPI-stained cells. Panels: (a) *cdc28-P8* at 25°C; (b) *cdc28-P8* after 4 h at 37°C; (c) *cdc28-P8* after 8 h at 37°C; and (d) wild-type cells at 37°C. Size bar, 5 μ M.

tested them for co-instability of leucine prototrophy and temperature-resistant (ts^+) growth. Plasmids from four ts^+ transformants were able to complement the ts^- growth defect when reintroduced into SP968. Restriction analysis of these plasmids showed that all four contained identical *Hind*III fragments with a total insert size of 7.5 kbp. One of these plasmids (pcdc28-1) was chosen for further analysis (Figure 2). Integration mapping confirmed that the isolated DNA fragment is the *cdc28*⁺ gene. Genomic mapping by hybridization to a cosmid library (Mizukami *et al.*, 1993) showed that *cdc28* lies between *cen2* and *ran1/pat1* on cosmid 1534 on the long arm of chromosome II.

A 4.8-kb *Bam*HI–*Bgl*II fragment (containing ~1 kb of pWH5 vector; Figure 2) retained the ability to complement the *cdc28-P8* mutation. We subjected this fragment to sequence analysis (Figure 3). The sequence contained an ORF of 2.9 kb that has the potential to encode a polypeptide of 968 amino acids with a predicted molecular mass of 111,500 Da. A comparison of this amino acid sequence with those in the databases by the BLAST program (Altschul *et al.*, 1990) revealed a similarity to a group of putative RNA-dependent ATPases or RNA helicases (Figures 3 and 4). The closest matches to the *cdc28*⁺ ORF were the Prp2, Prp16, and Prp22 proteins from *S. cerevisiae* (Burgess *et al.*, 1990; Chen and Lin, 1990; Company *et al.*, 1991). The *cdc28*⁺ gene product shares 56% identity with Prp2p and 50% identity with Prp22p and Prp16p in the central to carboxy-terminal region (580 amino acids) of each protein, which includes the DEAH box. The product of the *S. pombe* *prh1*⁺ gene (Inoue *et al.*, 1992) also shares 43% identity with Cdc28 in this region (Figure 4).

The *cdc28-P8* Strain Exhibits a Splicing Defect

The Prp2, Prp16, and Prp22 proteins each perform a distinct ATP-dependent step in pre-mRNA splicing (Company *et al.*, 1991; Schwer and Guthrie, 1991; Kim and Lin, 1993). We therefore tested for a general splicing defect in *cdc28-P8* by assaying the splicing of the U6 RNA precursor (pre-U6 RNA). U6 RNA is one of the spliceosomal small nuclear RNAs (snRNAs). The *S. pombe* U6 gene is interrupted by an intron whose structure is typical of nuclear pre-mRNA introns (Tani and Ohshima, 1989). In *S. pombe* pre-mRNA splicing (*prp*) mutants, the pattern of unspliced pre-U6 RNA accumulation is identical to that of pre-mRNAs (Pot-

ashkin and Frendewey, 1989). We compared the efficiency of pre-U6 RNA splicing in *cdc28-P8* with the wild type and with two *S. pombe* *prp* mutants: *prp1-1* (Potashkin *et al.*, 1989) and *prp5-1* (Kim and Frendewey, unpublished data). Total RNA was isolated from wild-type (972), *cdc28-P8* (SP968), *prp1-1* (JP23), and *prp5-1* (DK633) strains at 25°C and at 2, 6, and 8 h after the cells were shifted to 37°C. RNA from each strain was analyzed by Northern blotting (Figure 5). Two oligonucleotide probes that recognize the intron or second exon of pre-U6 RNA were used for hybridization.

Figure 5A shows the results of hybridization with the intron probe. The blot was autoradiographed for an extended period of time to allow for detection of rare or unstable intermediates or products of the splicing reaction (discussed below). Under these conditions, a band corresponding to the unspliced pre-U6 RNA was observed in nearly every lane. Essentially equal amounts of unspliced pre-U6 RNA accumulated in the wild-type and mutant strains at 25°C. The detection of U6 precursor at the permissive temperature was consistent with previous results (Potashkin and Frendewey, 1989), which indicated that the efficiency of splicing is reduced at lower temperatures. After shifting to 37°C, the pre-U6 RNA gradually disappeared in the wild type (972); none was detected 8 h after transfer to the restrictive temperature. In contrast, accumulation of unspliced precursor increased at 37°C in the splicing mutants and in *cdc28-P8* relative to the wild type and to the pre-U6 RNA levels at 25°C. A minor band visible below the pre-U6 RNA signal in the *cdc28* RNA after 6 h at 37°C could suggest accumulation of a splicing intermediate, but it was not produced in sufficient amounts for identification. Other faint bands seem to be nonspecific background, as they are variable and do not correlate with strain or temperature differences. The accumulation of pre-U6 RNA in the mutants persists for at least 8 h at 37°C. These results were confirmed by hybridization with a probe against U6 exon-2. The autoradiogram shown in Figure 5B (a shorter exposure than in 5A) demonstrates that, like *prp1-1* and *prp5-1*, *cdc28-P8* accumulates pre-U6 RNA at the restrictive temperature, whereas the wild type does not. The highly stable, mature U6 RNA persists at the restrictive temperature even in severe mutants such as *prp1*, which has a complete block in splicing (Potashkin and Frendewey,




Figure 2. Restriction map of the 4.8-kb complementing fragment. The hatched area represents vector sequence; the arrow indicates the 2.9-kb open reading frame (ORF).

AAAT AGCTTCATG CTTTTGGTG ATATTATTCT GGTTCGCCCT ACAAATGTC TTTAGAGCAA TATGTGCTG ACAAAGCGAT -111
 TTCGCTACTG GGAATGTCGG AACCTTCGCT GGTAGAATAC CTAATTGCAG AAGCGAAGGG CTCTTCAAGT TCTAATAACC TGTATCAGAA ACTCGTCAGT TTTGGTATGG -1

ATGGTGATGATCTCGCGTTAAAGAATTTGCTCATACGCTTTATGCTAGAAATCCAGAGAAAGGCTCAAGACCAGAAAGAACTACAATGCAAGAAAGAAAGAAAGAAACAGGGGATT 120
 M V M I L R L K N L L I R F M L E Y P E K A Q D R K K T M Q E R K K K E Q G I 40

TTACAAATGGAGCGTTTGAATTCAGGTTATGATTTGTTGATAGAACCTCAATCTCATGAGCGCCTGGGAAGCCTTTTGAAGAAAAAAGCCGTTCTAAAACCTCTAAGCGAGAAATAGCC 240
 L Q M E R L N S S Y D L L I E P Q S H E T P G K P L K K K S R S K T P K R E I A 80

AGGAGACAACGAGATGAGGATGAGTGGGAGAGTGACGAGTATGAGGAGGTATGGATGGAAGTGTAGTATCTCTATAGAAGAAGATTCAGTTTCTACCGATTCTCAAATCATGATTAT 360
 R R Q R D E D E W E S D E Y E E V V D G S A S H P I E E D S V S T D F Q N H D Y 120

GAAAAAGCTCAGATCTGAGACAGAAAGACTCAATGATTTACGTGAAAGAGAAGAAATTTGAAGCGCTCTTAGACGTAAGACCTTGAAGCCGCAACAAATGAATTTTGGAGGATTAT 480
 E K S S D P E T E R L N D L R E R E E F E E R L R R K D L E A A T N E F V E D Y 160

TCATCCAAATTTTCTCGGAAGAGCTAGCTCTTAGAAAACAGCAGATGATCTGAATCATGGAAGAAACTTGCTTCTGAACTTCGAAAAAATCGCGGCAACAATATCTTAAACCTAGG 600
 S S K F S S E E L A L R K L A D D P E S W R K L A S E L R K K S R Q Q Y L K P R 200

GCTCAGCAACAGTTGGAATCTTCCGAGAGAAATTCGAGATGAGGAGCAACTTTTTCAGGCGAGAAAGTTGACGCAAGCAGAAATAGAGAGCTTGAAGAAAAAAGAACTACTTTCGA 720
 A Q Q Q L E I L R R E I R D E E Q L F A G E K L T Q A E I R E L E K K K E L L R 240

ATTGCTGAAGAACGTCAGAGATTGAAAAACAAGCCACTGAATATCAAATGCCTGAAGACTATTTTACAGAGCAAGAAAATTTGACAGAAAAAGAAAGAGGAGTCTTATATCAAAGG 840
 I A E E R Q R L E K Q A T E Y Q M P E D Y F T E Q G K L D R K R K E E V L Y Q R 280

TATAAGGATAGTAAATGAAGGTGAGCAAAACGAGGTTACGATGGGAGCTGCCAACAACAGCGTGGGAGCTCAACAAATAAATAAGCAATTAATTTTATGATCAAAACGAATGGCTTCCT 960
 Y K D S N E G E Q N E V T M G A A E Q Q R W E A Q Q I N K A L L F D Q N E W L P 320

CCAGGAGAGAAGCAATTTGATTTCTGATTTGACGAAATCGCAACAGATTGATTTTCTACTCGATCTAAGCTTTCTGCGGAGAACTCTGTGACACTGCAAGATGACTGATGTTAAAGTT 1080
 P G E K Q F D F V F D E S Q Q I D F L L D T K L S A E N P V D T D K M T D V K V 360

GAAAAATCATTAGAAAGTTCTCGAAAAAGTTTACCTGTTTATCAATACAAGGACGATCTTTTGAAGCAATAAATGAGTACCAAGTGTGCTTATTTGTCGCTGAGACCGGATCTGTTGATAA 1200
 E K S L E S S R K S L P V Y Q Y K D D L L K A I N E Y Q V L L I V A E T G S G K 400

ACAACCTCAACTTCTCAATTTCTGCACGAAGCTGTTTATACTAAAGGAAATTAAGAAAATTTGCTGCACCAACCAAGAGAGTGTGCTGCAATGCTGTTGCTGCTAGGTTGCCAAGGAA 1320
T T Q L P Q F L H E A G Y T K G N K K I C C T O P R R V A A M S V A A R V A K E 440

ATGGATGTCAGGCTTGGACAGAAAGTTGGGTACAGCATTCGTTTGAATAATGCTACTTCTGAAAAAATGTCATAAAATACTGACTGATGGCATGCTTCTCAGAGAAATTTTAAACAGAA 1440
 M D V R L G Q E T E R Y S I R F E N A T S E K T V I K Y L T D G M L L R E F L T E 480

CCTGACCTGGCAAGCTATTCTGTCTATTATTATAGATGAAGCCATGAAGAAGCCTTCATACAGACATTTTATTGGACTTGTAAAGGATATTGCTAGATTTCGTCAGATTTAAAGGTT 1560
 P D L A S Y S V I I I D E A H E R T L H T D I L F G L V K D I A R F R P D L K V 520

CTAATTTGAGTGTACTACTATAGATGCCGAAAAATTTTCGGCTTACTTTGATGAAGTCCAGTGTCTATGTTCTGCGAGAAGATCCAGTGCATTTTATTACACCCCGCAACCTGAA 1680
 L I S S A T I D A E K F S A Y F D E A P V F Y V P G R R Y P V D I Y Y T P Q P E 560

GCAAAATATATACAAGCGCTATTACAACCTTCTTCAAAATACATACTACCAGCCTGCTGGTATATTTAGTTTTTTTAAACGGGCAAGATGAAATGAAATTAATGTCGAAAAACATG 1800
 A N Y I Q A A I T T I L Q I H T T Q P A G D I L V F L T G Q D E I E L M S E N M 600

CAAGAATTTGTCGGATACTAGGAAAAAGAAATTCGGAAATAATTTTATGTCGATTTATGCCAATCTGCCCTTGAATGCAAGCTAAAAATTTTATGATCTACACCCCTGGTGCCTCCG 1920
 Q E L C R I L G K R I P E I I L C P I Y A N L P S E L Q A K I F D P T P P G A R 640

AAAGTTGATTAAGCAACCAATATTGCTGAAACTTCTATAACAATGACGAGTGAATTTTGTGATGATTTGATTTGCAAAACAAACATGATAATCAAGGACTGGTATGGAAGC 2040
 K V V L A T N I A E T S I T I D G V N F V I D S G F V K Q N M Y N P R T G M E S 680

TTAGTATCTGTGCTTCTCCCGTCTCTGCTGATCAGCGTCCCGCGCTGCTGGTGGAGTAGGACCTGGAAGTGTCTCCGTTTATATACCGCTTGGACATATAACAATGAAGTGGAT 2160
 L V S V P C S R A S A D Q R A G R A G R V G P G K C F R L Y T R W T Y N N E L D 720

ATGGTAAACATCTCCTGAAATTCAAAGAACAAATTTGACAAATATGTTCTTCTGCTTAAATCCCTAGGAATTAATAATCTTTTATGATTTGACTTCAATGATGCTCCACCCTGAAACC 2280
 M V T S P E I Q R T N L T N I V L L L K S L G I N N L L D F D F M D A P P P E T 760

TTGATGCGTTCAATGAGTATTGATGCGCTTGGTCCCTTAAACAATAGAGGAGAAATGACTAAGCTCGGAGCAGATGGCAGAGTTTCTCAGATCTATGCTTTTCTAAATCTTTG 2400
 L M R S L E L L Y A L G A L N N R G E L T K L G R Q M A E F P T D P M L S K S L 800

ATTGCATCTTCAAGATGAGTGTGTTGAAGAAGTGTATCAATGTTTCTATGTTAGGGGAGGCTTCTCTTTTATCTATAGACAAAAGACAAAATATGGAAGCGGATAAAGCTAGA 2520
 I A S S K Y G C V E E V L S I V S M L G E A S S L F Y R P K D K I M E A D K A R 840

GCGAATTTACCCAACTGGTGGTATCAATTTAACTCTTCTCATATTTGGAATGAATGGGTAGATGACTTTTCTGATCAATTTGGCCCGTGAAGAACTTTTACAGTATAAGAGTTTA 2640
 A N F T Q P G G D H L T L L H I W N E W V D T D F S Y N W A R E N F L Q Y K S L 880

TGTCGAGCTCAGAGATGTTGATCAACTTGCCTTCTGTAACCGGTGAAATTTGAATTTGTAATGTAATTTCTCAGAGTCCCTTGAATCTTAAAGAAAGCTATCAGAGCTGGTTAT 2760
 C R A R D V R D Q L A N L C E R V E I E L V T N S S E S L D P I K K A I T A G Y 920

TTTCTAATGCTGACGCTTGGATGCTGATGGGATTTCTTATGCACCTGTTAAAGTAATCAAACCGTTTATATTCACCCCTAGTCTTCTGATGCTGAGAAAAAGCCAAAGTCAATCATT 2880
 F P N A A R L D R S G D S Y R T V K S N Q T V Y I H P S S V A E K K P K V I I 960

TACTTTGAATGGGTTTGGACAAC TAA GGAGTATTGC AGACAAATTA CTGAAATACA ACCCGAGTGG CTCTTGGAAA TAAGCCCCA TTATTTCAAG CCAGAAAAA 2987
 Y F E L G F D N . 968

TAGAAGAAIT GCAAAGACG CAAAACGAC ATAAGCGSTA AAAGAGTGA AAAGGAAAT TAGTTTTAAA GGCGATTTGG TTGTATATAA TCATAATGAA TTTATGTTTC 3097
 ACAATAATGG ACAAACTTTT AAAAGATCCC CCGGG 3132

Figure 3.

	I	Ia	
cdc28	VLLIVAETGSGKTTQLPQFLHEAGYTKGNK-KICCTOPRRVAAMSVAARVAKEMDVRVGQEVGYSIRFENAT-SEKTVIKYLT		469
prh1	VIVVVGETGSGKSTQIPQFLNECPYAQEG--CVAITOPRRVAAVNLAARVAEQGCRLEQVGYSIREFDDTT-SKKTRIKYLT		193
PRP22	FLVIVGETGSGKTTQITQYLDEEGFSNYG--MIGCTOPRRVAASVAARVAEEVGCKVGHVGYTIRFEDVT-GPDTRIKYMT		580
PRP16	VVVIIGETGSGKTTQLAQYLYEEGYANDRGKSI VVTOPRRVAASVAARVAEMQVPLGKEVGYSIREFEDVTDSECTKLFVVT		450
PRP2	VLIIMGETGSGKTTQLPQYLVEDGFTDQGKLOIAITOPRRVAASVAARVADEMNVVLGKEVGYQIRFEDKTFPNTKTVLKYMT		323
	II	III	
cdc28	DGMLLREFLTEPDLASYSVIIIDEAHERTLHTDILFGLVKDIARFRPDLKVLISSATIDAEEKFSAYFDEAPVFYVPGRRYPVDIY		554
prh1	DGMLLRELINDPILSQYHTLLIDEAHERTLMTDMLLGFVVKI IKKRALRVIMSATLNAERFSEFFDGAEICYISGRQYFVQIH		278
PRP22	DGMLQREALLDPEMSKYSVIMLDEAHERTVATDVLFALLKKAAIKRPDLKVI VTSATLNSAKFSEYFLNCP I INIPGKTFPVEVL		665
PRP16	DGILLRETLDDTLDKYS CVIIDEAHERSLNTDILLGFFKILLARRDLKLIIT SATMNAKFSAFFGNAPQFTIPGRTFPVQTI		535
PRP2	DGMLLREFLTDKSLKYS CIMIDEAHERTLATDILIGLLKDI LPQRP TLKLLISSATMNAKFS EFFDNCP IFNVPGRRYPVDI H		408
	IV		
cdc28	YTPQPEANYIQAAITITLQOIH TTQPA--GDILVFLTGQDEIE LMSENMQELCRILGKRIP EI-----ILCPIYANLPSEL		628
prh1	YTYTPEPDYLDACLRTIF-QLHTKLP--GDILVFLTGQDEIEALEALIKSYKQLPSNLPQI-----QACPLFASLPQEQ		352
PRP22	YSQTPQMDYIEAALDCVI-DI H INEGP--GDILVFLTGQDEIE DSCCEILYDRVKT LGDSIGEL-----LILPVYSALPSEI		739
PRP16	YTSNPVQDYVEAAVSQAV-KIHLANDCSSGDILIFMTGQEDIE TTFDTLQEKFLQVYSKFKGTANFEEINDIEILPIYSALPADL		620
PRP2	YTLQPEANYIHAAITTF-QIHTTQSLP-GDILVFLTGQDEIE IERTKT----KLEEIMSKLGSRTKQM-----ITPIYANLPQEQ		483
	V	VI	
cdc28	QAKIFDPTPPGARKVVLATNIAETSITIDGVNFFVIDSGFVKQNMYPRTGMESLVSVP CSRASADQRAGRGRVGP GKCFRLYT		711
prh1	QLQVFLPALANHRKVVLSSTNIAETS SVTISGIRYVIDTGLAKIKQFNSKLGLES LTVPISQSAAMQRSGRAGREAAQC YRIYT		435
PRP22	QSKIFEP TPKGRSRKVV FATNIAETSITIDGIYVVDPGFAKINIYNARAGIEQLIVSPI SQAQANQRKGRAGRTGPGKCYRLYT		822
PRP16	QFKIFQDLHGTRKRI I IATNIAETSLTIKGRYVIDCGYSKLVKNPKIGLDSLVI TPI SKANADQRSGRAGR TAPGTRYRLYT		703
PRP2	QLKIFQPTPENCRKVV L ATNIAETSLTIDGIRYVIDPGFVKENSYPVSTGMTQLLTVPCSRASVDQRAGRGRVGP GKCFRIFT		566
cdc28	RWTYNNELDMVTSPEIQRTNL TNIVLLKSLGINNLL-DFDFMDAPP PETLMRSLELLYALGALNNRGELTKLGRQMAEFP TDPML		796
prh1	-EADFDKLPKETIPEIKRIDLSQAVLTLKARGQNDVI-NFHYMDPPSKEGLLRALHELYSIGALDDNGHINDLGYQMSLIP LLLPSL		520
PRP22	ESAFYNEMLNTVPEIQRONLSHTI LMLKAMGINDLL-KFDFMDPPPKNMLNALTELYHLQSLDDEGKLTNLGKEMSLFPMDPTL		907
PRP16	EDTFKEDMYLQTIPEIQRTNLSNTLLKSLDVTDELSKFF IDKPLQTF LSSLYELWFI GAIDTSGQLTFLGLQMAKFFLPQSL		789
PRP2	KWSYLHELELMPKPEITRTNLSNTVLLLSLGVTDLI-KFPLMDKPSIPTLRKSLNLYILGALNSKGTITRLGKMMCEFPCEPEF		651
cdc28	SKSLIASSKYG-C---VEEVL SIVSMLGEASSLFYRPKDKIMEADKARANFTQPGGDHLLTLLHIWNEWVD TDFSYNWARENFLOQYK		878
prh1	ARAVLAAREHN-C---LSEVIDVVSCLSTDSMFLFPQEKR-DEAIEARLKFHLHSEGDLTCLNALRQYLES HSDSRKQWCSQNF IN		601
PRP22	SRSLLSSVDNQ-C---SDEIVTII SMLSVQNVF YRPKDRQ-LEADSKAKFHHPYGDHLLTLLNVYTRWQYANYSEQYCKTNFLHFR		988
PRP16	SKILLIAVRNG-C---SDEMLTIVSMLSVPQVYRPKERQ-KEAD IARNKFF IAKSDHLLTLLNVFEQWRANNFSSHWCNKHFVQYK		870
PRP2	AKVLYTAATHEQCQGVLEECLTIVSMLHETPSLFIG----QKRDAASVLSVEVSDHLLYLEIFNQWRNSKFSRSWCQDHKIQFK		732

Figure 4. Alignment of the central region of the Cdc28 protein sequence with similar members of the DEAH-box RNA helicase family. The Prh1 protein is from *S. pombe*, and the Prp2, Prp16, and Prp22 proteins are from *S. cerevisiae*. Amino acids that are identical in all five or four of five of the proteins are emphasized in bold lettering. Roman numerals indicate the position of the motifs found in the DEAH-box putative RNA helicases (Company *et al.*, 1991).

1989). These results establish that *cdc28-P8* exhibits a general defect in splicing roughly equivalent to that seen in two severe splicing mutants.

To confirm that *cdc28-P8* has a defect in pre-mRNA splicing as well as in pre-U6 RNA splicing, we performed Northern blot analysis with the use of probes for the TFIID mRNA. The *S. pombe* TFIID gene has

three introns (Hoffmann *et al.*, 1990). As shown in Figure 6, the amount of the unspliced TFIID pre-mRNA greatly increases in *cdc28-P8* (lane 6), as it does in the two pre-mRNA splicing mutants *prp2-2* and *prp8-1* (lanes 4 and 8), after shifting to the restrictive temperature for 2 h (lane 6). In contrast, the wild-type cells do not accumulate pre-mRNA under the same conditions (lane 2). There is also a reduction in mature mRNA levels of the mutants at the restrictive temperature, which indicates a nearly complete block in splicing. These results establish that *cdc28-P8* exhibits a general defect in splicing roughly equivalent to that seen in four severe splicing mutants.

Figure 3 (cont). Nucleotide sequence of the 2904-bp ORF in the *cdc28+* gene and its predicted amino acid sequence (accession number U48733). No introns were detected. Restriction sites are indicated and underlined. The seven conserved regions for DEAH-box RNA helicases are underlined.




Figure 5. Northern blot analysis of pre-U6 RNA splicing. Wild-type (972) and mutant strains (*cdc28*, *prp1*, and *prp5*) were grown to mid-log phase and shifted to 37°C. Total RNA was isolated at 0, 2, 6, and 8 h after the temperature shift, analyzed on a 10% polyacrylamide/7 M urea gel, and transferred to Hybond membrane. The blot was probed with ³²P-labeled oligonucleotides U6 I-2 (A) and U6 E2-2 (B). Pre-RNA (P) is 149 nucleotides, and spliced, mature U6 RNA (M) is 99 nucleotides. The sizes in nucleotides of DNA markers are indicated on the left in each panel.

cdc28 Is Allelic with *prp8*

Because *cdc28-P8* has a temperature-sensitive splicing defect, we wanted to determine whether *cdc28*⁺ was identical to any of the genes identified by mutations that affect pre-mRNA splicing. Crosses between




Figure 6. Northern blot analysis of pre-mRNA splicing. Wild-type (972) and mutant strains (*cdc28-P8*, *prp2-2*, and *prp8-1*) were grown to mid-log phase at 26°C and either maintained at 26°C or shifted to 36°C for 2 h. Total RNA was isolated from each strain and then subjected to Northern blot analysis. The blot was hybridized with a mixture of ³²P-labeled oligodeoxynucleotides TFII-IN1 and TFII-EX3, which are complementary to the first intron and the third exon of the pre-TFIID mRNA, respectively. The RNA source strains and the incubation temperatures are indicated above the lanes. WT represents the wild-type cells. Top and bottom arrowheads on the right indicate the positions of the unspliced precursor (pre) and mature (m) TFIID mRNAs.

cdc28-P8 and the previously described *ts*⁻ alleles *prp1-1*, *prp2-1*, *prp3-1* (Potashkin *et al.*, 1989), and *prp4-1* (Rosenberg *et al.*, 1991) or the new alleles *prp5-1*, *prp6-1*, *prp7-1*, and *prp9-1* (Kim and Frendewey, unpublished data) produced diploid progeny that were able to grow at 37°C. However, diploids produced from a cross between *cdc28-P8* and a new splicing mutant *prp8-1* remained temperature sensitive. Thus, *cdc28* and *prp8* are in the same complementation group. Genetic linkage analysis corroborated this finding. We found no temperature-resistant progeny among ~1400 random spores from a cross of *cdc28-P8* with *prp8-1*. From this tight linkage and the complementation grouping we conclude that *cdc28* and *prp8* are identical loci.

cdc Phenotype of *prp8-1*

The *prp8-1* mutant divided twice before it ceased growing at the restrictive temperature. Like *cdc28-P8*, the cells began to elongate 6 h after the shift to the restrictive temperature. Almost all of the *prp8-1* cells elongated two- to threefold compared with the wild-type cells after 8 h at the restrictive temperature. Elongation of the cells at the restrictive temperature is a typical phenotype of the *cdc* mutants. Although the nuclear material in *cdc28-P8* seemed to be condensed after an 8-h shift, *prp8-1* did not show such a phenotype under the same conditions. Thus, chromatin condensation may be an allele-specific phenotype of *cdc28-P8*.

DISCUSSION

The *S. pombe cdc28-P8* strain was isolated in a screen for *cdc* mutants and was found to be blocked in late G2 or entry into mitosis at the restrictive temperature (Nasmyth and Nurse, 1981). We have confirmed the G2 arrest in *cdc28-P8* and showed that it is followed by a rapid decrease in cell viability. When we cloned the *cdc28⁺* gene by complementation of the temperature-sensitive growth defect in *cdc28-P8*, we were surprised to find that it encoded a protein of 111-kDa—predicted molecular mass that is similar to a family of putative RNA-dependent ATPases or helicases required for pre-mRNA splicing in budding yeast. Consistent with this similarity, *cdc28-P8* exhibits a block in the splicing of the pre-mRNA-type intron in the pre-U6 RNA (Tani and Ohshima, 1989) as well as in the splicing of pre-mRNAs at the restrictive temperature. An independent screen for *S. pombe* mutants defective in pre-mRNA splicing identified the *prp8-1* strain that, in addition to its splicing defect, also expresses a *cdc* phenotype seemingly identical to that of *cdc28-P8*. We demonstrated that *cdc28* and *prp8* are identical loci. Thus, a mutation in a single gene can cause a specific arrest in progression through the cell cycle and a block in pre-mRNA splicing.

The *cdc28⁺/prp8⁺* gene product is most similar to the Prp2, Prp16, and Prp22 proteins of *S. cerevisiae*, which are members of the DEAH-box subfamily of putative RNA helicases. The product of the *S. pombe prh1⁺* gene (Inoue *et al.*, 1992), whose function is not known, is also a member of this family. Other possible members of the DEAH-box family have been identified in animal viruses (Koonin and Senkevich, 1992; Yanez *et al.*, 1993), humans (Ono *et al.*, 1994), budding yeast (James *et al.*, 1994), and *Escherichia coli* (Moriya *et al.*, 1995). The Cdc28/Prp8 protein is ~50% identical to Prp2p, Prp16p, and Prp22p in the most highly conserved C terminus. The eight characteristic motifs within the DEAD-box family of potential RNA helicases (Schmid and Linder, 1992) that are retained in the DEAH subfamily (Company *et al.*, 1991) are also present in Cdc28/Prp8, including the DEAH signature (amino acids 492–495), and TQPPRVAA (amino acids 423–430) and QRAGRAGR (amino acids 693–700), which are important for helicase activity (Company *et al.*, 1991; Schmid and Linder, 1992).

The Prp2, Prp16, and Prp22 proteins act at three sequential steps in pre-mRNA splicing. Prp2p is required early, just before the first transesterification reaction that results in cleavage at the 5' splice site to produce the bimolecular intermediate, which consists of the upstream exon and the downstream exon attached to the intron in a lariat form (Lin *et al.*, 1987; King and Beggs, 1990; Kim and Lin, 1993). The second transesterification reaction that produces the spliced exons and releases the intron lariat requires Prp16p

(Schwer and Guthrie, 1991). The final step in pre-mRNA splicing—the release of the spliced mRNA and intron lariat from the spliceosome—is promoted by Prp22p (Company *et al.*, 1991). These functions were first suggested by the molecules that accumulate in mutant strains: unspliced pre-mRNA in *prp2*, intermediates in *prp16*, and intron lariat in *prp22*. The *cdc28-P8* and *prp8-1* mutants accumulate unspliced precursors. Although a faint band was observed, splicing intermediates or final products were not observed to accumulate to significant levels in the mutants. This phenotype supports the notion that *S. pombe* Cdc28/Prp8 is homologous to *S. cerevisiae* Prp2p, but co-complementation experiments and further functional studies will be required to confirm this hypothesis.

The expression of both *cdc* and *prp* phenotypes in *cdc28-P8* and *prp8-1* joins a growing list of observations that point to a connection between RNA metabolism and regulation of the cell-division cycle. The earliest reported example of this phenomenon may be the peculiar observations that changes in tRNA expression can influence mitosis in fission yeast (Nurse and Thuriaux, 1984). Similar observations have recently been made in budding yeast (Reijo *et al.*, 1993). A better-understood example is the abundant mammalian nuclear protein RCC1, which prevents premature initiation of mitosis (Nishimoto *et al.*, 1978; Dasso, 1993). RCC1 is the guanine nucleotide exchange factor for the Ran GTPase, a central regulator of nuclear-cytoplasmic trafficking (Rush *et al.*, 1996). Mutations in the *S. cerevisiae* homologue of RCC1, *PRP20/SRM1*, have pleiotropic effects on nuclear structure and RNA metabolism, including accumulation of nuclear poly(A)⁺ RNA and defects in pre-mRNA splicing (Aebi *et al.*, 1990). Two recent genetic results, whose meaning is at the moment obscure, support the notion of RNA-mediated events in the control of the cell cycle. In one case a *Drosophila* gene that encodes a DEAD-box RNA helicase can suppress loss of the key cell-cycle regulators Wee1 and Mik1 in *S. pombe* (Warbrick and Glover, 1994), and a gene that encodes a nuclear poly(A)⁺ RNA binding protein (Wilson *et al.*, 1994) was isolated as a suppressor of a dominant mutation in the *S. cerevisiae* *CLN3* gene (Sugimoto *et al.*, 1995), which encodes a G1 cyclin.

More relevant to *cdc28⁺*, the *dbf3-1* mutation in the *S. cerevisiae* *PRP8* gene, which encodes a protein of the U5 snRNP, produces a cell-cycle block before S phase (Shea *et al.*, 1994). Similar results were obtained with the budding-yeast mutants *dbf2* and *dbf5*, which also exhibit an S-phase arrest. *DBF5* was shown to be allelic with *PRP3* (Johnston and Thomas, 1982), and the *USS1* gene, which encodes a U6 snRNA binding protein, was isolated as a multicopy suppressor of *dbf2* (Cooper *et al.*, 1995). The *mis11-453* (Takahashi *et al.*, 1994) mutation, which results in minichromosome loss in *S. pombe*, was shown to reside in the *prp2⁺* gene, which

encodes a homologue of the large subunit of the human pre-mRNA splicing factor U2AF (Potashkin *et al.*, 1993). In a related connection between chromosome dynamics and pre-mRNA splicing, the *S. pombe* *dsk1*⁺ gene, which encodes a kinase, was isolated as a multicopy suppressor of *dis1* mutants, which fail to separate sister chromatids at mitosis (Ohkura *et al.*, 1988; Takeuchi and Yanagida, 1993). The human homologue of Dsk1, SRPK1, causes nuclear disruption when overexpressed (Gui *et al.*, 1994). The SRPK1 kinase specifically phosphorylates members of the SR family of pre-mRNA splicing factors (Fu, 1995). The *dis1*⁺ gene encodes a protein that associates with microtubules and the spindle pole bodies (Nabeshima *et al.*, 1995), a function that seems to have no connection with RNA processing. However, a conditional-lethal allele of *S. cerevisiae* PRP22 elicits a loss of microtubules and a *cdc* arrest that produces large-budded cells at the restrictive temperature (T. Huffaker, personal communication). These disparate observations, along with the variety of *cdc* defects that have been observed in newly isolated *S. pombe* *prp* mutants (Kim and Fren-dewey, unpublished results; S. Urushiyama, T. Tani, and Y. Ohshima, unpublished results), suggest an intricate interplay between RNA-processing events and the regulation of the cell cycle.

What mechanisms could explain how mutations in splicing factors affect regulation of the cell cycle? A simple proposal is that a block in the splicing of the pre-mRNAs for key cell-cycle regulators leads to cell-cycle arrest at a point at which the regulator is required. This model does not, however, satisfactorily explain why only some of the known *prp* alleles in both yeast strains produce *cdc* phenotypes. All of the known *prp* genes in both *S. pombe* and *S. cerevisiae* seem to be required for constitutive splicing of all pre-mRNAs, yet different *prp* mutants elicit different *cdc* defects. These observations suggest a more global explanation that takes into account the role of the pre-mRNA splicing apparatus in nuclear structure. The nucleus contains a complex ribonucleoprotein network, and pre-mRNA splicing factors are localized to organized sites within this substructure (Spector, 1993). Several lines of evidence suggest that the spliceosome (Zeitlin *et al.*, 1989) or some of its components (Blencowe *et al.*, 1994; Zeng *et al.*, 1994; Chabot *et al.*, 1995) reside on the nuclear matrix. It is interesting with respect to cell-cycle regulation that the retinoblastoma protein associates with a nuclear matrix protein at subnuclear regions actively involved in RNA metabolism (Durfee *et al.*, 1994). The SR protein-splicing factors (Fu, 1995), in particular, seem to be tightly associated with nuclear matrix proteins (Blencowe *et al.*, 1994). These interactions between the splicing apparatus and the nuclear scaffold may need to be disrupted or reorganized to allow other nuclear events, such as mitosis and DNA replication, to proceed. The

SRPK1/Dsk1 kinase (Ohkura *et al.*, 1988; Gui *et al.*, 1994) is a good candidate to promote this rearrangement of the ribonucleoprotein network of the nucleus. Likewise, RNA helicases, such as Prp22p from *S. cerevisiae* (Company *et al.*, 1991) and *cdc28/prp8* from *S. pombe*, may facilitate disruption of the spliceosome as a prerequisite for nuclear division. Certain splicing factors could therefore serve dual roles as functional subunits of the spliceosome and as structural components of the nuclear matrix. Thus, mutations that affect pre-mRNA splicing could also prevent dynamic changes in nuclear structure required for cell-cycle progression. Some splicing factors, for example the helicases, could provide independent functions in mitotic processes such as spindle formation or chromosome separation. This hypothesis raises the intriguing possibility that RNA-mediated events could be crucial to the regulation of cell division.

ACKNOWLEDGMENTS

We wish to thank Nancy Kaplan, Craig Gawel, and R. Massey for technical assistance and George Stephano for encouragement. Maureen Caligiuri, Dave Casso, Harriet Fielotter, and Greg Hannon are thanked for critical comments on this manuscript. We also thank Jim Duffy, Phil Renna, and Mike Ockler for artistic assistance. This work was supported by National Institutes of Health grant GM-34607 to D.B. and an American Cancer Society postdoctoral fellowship to K.L. D.B. is an investigator for the Howard Hughes Medical Institute. D.F. is the recipient of an Irma T. Hirsch Career Scientist Award.

REFERENCES

- Aebi, M., Clark, M.W., Vijayraghavan, U., and Abelson, J. (1990). A yeast mutant, *PRP20*, altered in mRNA metabolism and maintenance of nuclear structure, is defective in a gene homologous to the human gene *RCC1*, which is involved in the control of chromosome condensation. *Mol. Gen. Genet.* 224, 72–80.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local alignment search tool. *J. Mol. Biol.* 215, 403–410.
- Aves, S.J., Durkacz, B.W., Carr, A., and Nurse, P. (1985). Cloning, sequencing, and transcriptional control of the *Schizosaccharomyces pombe* *cdc10* "start" gene. *EMBO J.* 4, 457–463.
- Beach, D., Piper, M., and Nurse, P. (1982). Construction of a *Schizosaccharomyces pombe* gene bank in a yeast bacterial-shuttle vector and its use to isolate genes by complementation. *Mol. Gen. Genet.* 187, 326–329.
- Beach, D., Rodgers, L., and Gould, J. (1985). *RAN1*⁺ controls the transition from mitotic division to meiosis in fission yeast. *Curr. Genet.* 10, 297–311.
- Blencowe, B.J., Nickerson, J.A., Issner, R., Penman, S., and Sharp, P.A. (1994). Association of nuclear matrix antigens with exon-containing splicing complexes. *J. Cell Biol.* 127, 593–607.
- Booher, R., and Beach, D. (1987). Interaction between *cdc13*⁺ and *cdc2*⁺ in the control of mitosis in fission yeast: dissociation of the G1 and G2 roles of the *cdc2*⁺ protein kinase. *EMBO J.* 6, 3441–3447.
- Booher, R., and Beach, D. (1988). Involvement of *cdc13*⁺ in mitotic control in *Schizosaccharomyces pombe*: possible interaction of the gene product with microtubules. *EMBO J.* 7, 2321–2327.

- Bueno, A., Richardson, H., Reed, S.I., and Russell, P. (1991). A fission-yeast B-type cyclin functioning early in the cell cycle. *Cell* 66, 149–159.
- Bueno, A., and Russell, P. (1993). Two fission-yeast B-type cyclins, Cig2 and Cdc13, have different functions in mitosis. *Mol. Cell. Biol.* 13, 2286–2297.
- Burgess, S., Couto, J.R., and Guthrie, C. (1990). A putative ATP binding protein influences the fidelity of branch point recognition in yeast splicing. *Cell* 60, 705–717.
- Caligiuri, M., and Beach, D. (1993). Sct1 functions in partnership with Cdc10 in a transcription complex that activates cell-cycle START and inhibits differentiation. *Cell* 72, 607–619.
- Chabot, B., Bisotto, S., and Vincent, M. (1995). The nuclear matrix phosphoprotein p255 associates with splicing complexes as part of the [U4/U6.U5] tri-snRNP particle. *Nucleic Acids Res.* 23, 3206–3213.
- Chen, J.H., and Lin, R.J. (1990). The yeast PRP2 protein, a putative RNA-dependent ATPase, shares extensive sequence homology with two other pre-mRNA splicing factors. *Nucleic Acids Res.* 18, 6447.
- Company, M., Arenas, J., and Abelson, J. (1991). Requirement of the RNA helicase-like protein PRP22 for release of messenger RNA from spliceosomes. *Nature* 349, 487–493.
- Connolly, T., and Beach, D. (1994). Interaction between the Cig1 and Cig2 B-type cyclins in the fission-yeast cell cycle. *Mol. Cell. Biol.* 14, 768–776.
- Cooper, M., Johnston, L.H., and Beggs, J.D. (1995). Identification and characterization of Uss1p (Sdb23p): a novel U6 snRNA-associated protein with significant similarity to core proteins of small nuclear ribonucleoproteins. *EMBO J.* 14, 2066–2075.
- Costello, G., Rodgers, L., and Beach, D. (1986). Fission yeast enters the stationary phase G0 state from either mitotic G1 or G2. *Curr. Genet.* 11, 119–125.
- Dasso, M. (1993). RCC1 in the cell cycle: the regulator of chromosome condensation takes on new roles. *Trends Biochem. Sci.* 18, 96–101.
- Draetta, G. (1990). Cell-cycle control in eukaryotes: molecular mechanisms of cdc2 activation. *Trends Biochem. Sci.* 15, 378–383.
- Draetta, G., Piwnicka-Worms, H., Morrison, D., Druker, B., Roberts, T., and Beach, D. (1988). Human cdc2 protein kinase is a major cell cycle-regulated tyrosine kinase substrate. *Nature* 336, 738–744.
- Ducommun, B., Brambilla, P., Felix, M.A., Franza, B., Jr., Karsenti, E., and Draetta, G. (1991). cdc2 phosphorylation is required for its interaction with cyclin. *EMBO J.* 10, 3311–3319.
- Durfee, T., Mancini, M.A., Jones, D., Elledge, S.J., and Lee, W.H. (1994). The amino-terminal region of the retinoblastoma gene product binds a novel nuclear matrix protein that colocalizes to centers for RNA processing. *J. Cell Biol.* 127, 609–622.
- Fernandez Sarabia, M.J., McNerny, C., Harris, P., Gordon, C., and Fantes, P. (1993). The cell-cycle genes *cdc22⁺* and *suc22⁺* of the fission yeast *Schizosaccharomyces pombe* encode the large and small subunits of ribonucleotide reductase. *Mol. Gen. Genet.* 238, 241–251.
- Forsburg, S.L., and Nurse, P. (1991). Identification of a G1-type cyclin *puc1⁺* in the fission yeast *Schizosaccharomyces pombe*. *Nature* 351, 245–248.
- Fu, X.D. (1995). The superfamily of arginine/serine-rich splicing factors. *RNA* 1, 663–680.
- Gould, K.L., Moreno, S., Owen, D.J., Sazer, S., and Nurse, P. (1991). Phosphorylation at Thr167 is required for *Schizosaccharomyces pombe* p34^{cdc2} function. *EMBO J.* 10, 3297–3309.
- Gui, J.F., Lane, W.S., and Fu, X.D. (1994). A serine kinase regulates intracellular localization of splicing factors in the cell cycle. *Nature* 369, 678–682.
- Hartwell, L.H. (1978). Cell division from a genetic perspective. *J. Cell Biol.* 77, 627–637.
- Henikoff, S. (1987). Unidirectional digestion with exonuclease III in DNA sequence analysis. *Methods Enzymol.* 155, 156–165.
- Hiraoka, Y., Toda, T., and Yanagida, M. (1984). The NDA3 gene of fission yeast encodes β -tubulin: a cold-sensitive *nda3* mutation reversibly blocks spindle formation and chromosome movement in mitosis. *Cell* 39, 349–358.
- Hoffmann, A., Horikoshi, M., Wang, C.K., Schroeder, S., Weil, P.A., and Roeder, R.G. (1990). Cloning of the *Schizosaccharomyces pombe* TFIID gene reveals a strong conservation of functional domains present in *Saccharomyces cerevisiae* TFIID. *Genes Dev.* 4, 1141–1148.
- Inoue, S.B., Sakamoto, H., Sawa, H., and Shimura, Y. (1992). Nucleotide sequence of a fission-yeast gene encoding the DEAH-box RNA helicase. *Nucleic Acids Res.* 20, 5841.
- Ito, H., Fukuda, Y., Murata, K., and Kimura, A. (1983). Transformation of intact yeast cells treated with alkali cations. *J. Bacteriol.* 153, 163–168.
- James, C.M., Gent, M.E., Indge, K.J., and Oliver, S.G. (1994). Sequence analysis of a 10-kb fragment of yeast chromosome XI identifies the *SMY1* locus and reveals sequences related to a pre-mRNA splicing factor and vacuolar ATPase subunit C plus a number of unidentified open reading frames. *Yeast* 10, 247–255.
- Johnston, L.H., and Thomas, A.P. (1982). The isolation of new DNA synthesis mutants in the yeast *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* 186, 439–444.
- Kim, S.H., and Lin, R.J. (1993). Pre-mRNA splicing within an assembled yeast spliceosome requires an RNA-dependent ATPase and ATP hydrolysis. *Proc. Natl. Acad. Sci. USA* 90, 888–892.
- King, D.S., and Beggs, J.D. (1990). Interactions of PRP2 protein with pre-mRNA splicing complexes in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 18, 6559–6564.
- Koonin, E.V., and Senkevich, T.G. (1992). Vaccinia virus encodes four putative DNA and/or RNA helicases distantly related to each other. *J. Gen. Virol.* 73, 989–993.
- Leupold, U. (1970). Genetical methods for *Schizosaccharomyces pombe*. *Methods Cell Physiol.* 4, 169–177.
- Lin, R.J., Lustig, A.J., and Abelson, J. (1987). Splicing of yeast nuclear pre-mRNA in vitro requires a functional 40S spliceosome and several extrinsic factors. *Genes Dev.* 1, 7–18.
- Lowndes, N.F., McNerny, C.J., Johnson, A.L., Fantes, P.A., and Johnston, L.H. (1992). Control of DNA synthesis genes in fission yeast by the cell-cycle gene *cdc10⁺*. *Nature* 355, 449–453.
- Lundgren, K., Walworth, N., Booher, R., Dembski, M., Kirschner, M., and Beach, D. (1991). *mik1* and *wee1* cooperate in the inhibitory tyrosine phosphorylation of cdc2. *Cell* 64, 1111–1122.
- Matsumoto, T., and Beach, D. (1991). Premature initiation of mitosis in yeast lacking RCC1 or an interacting GTPase. *Cell* 66, 347–360.
- Millar, J.B., Lenaers, G., and Russell, P. (1992). Pyp3 PTPase acts as a mitotic inducer in fission yeast. *EMBO J.* 11, 4933–4941.
- Mizukami, T., Chang, W.I., Garkavtsev, I., Kaplan, N., Lombardi, D., Matsumoto, T., Niwa, O., Kounosu, A., Yanagida, M., Marr, T.G., and Beach, D. (1993). A 13-kb resolution cosmid map of the 14-Mb fission-yeast genome by nonrandom sequence-tagged site mapping. *Cell* 73, 121–132.
- Moriya, H., Kasai, H., and Isono, K. (1995). Cloning and characterization of the hrpA gene in the terC region of *Escherichia coli* that is

- highly similar to the DEAH family RNA helicase genes of *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 23, 595–598.
- Morla, A.O., Draetta, G., Beach, D., and Wang, J.Y. (1989). Reversible tyrosine phosphorylation of *cdc2*: dephosphorylation accompanies activation during entry into mitosis. *Cell* 58, 193–203.
- Nabeshima, K., Kurooka, H., Takeuchi, M., Kinoshita, K., Nakaseko, Y., and Yanagida, M. (1995). *p93^{dis1}*, which is required for sister chromatid separation, is a novel microtubule and spindle pole body-associating protein phosphorylated at the Cdc2 target sites. *Genes Dev.* 9, 1572–1585.
- Nasmyth, K.A. (1977). Temperature-sensitive lethal mutants in the structural gene for DNA ligase in the yeast *Schizosaccharomyces pombe*. *Cell* 12, 1109–1120.
- Nasmyth, K., and Nurse, P. (1981). Cell-division cycle mutants altered in DNA replication and mitosis in the fission yeast *Schizosaccharomyces pombe*. *Mol. Gen. Genet.* 182, 119–124.
- Nishimoto, T., Eilen, E., and Basilico, C. (1978). Premature chromosome condensation in a *ts⁻* DNA mutant of BHK cells. *Cell* 15, 475–483.
- Nurse, P. (1975). Genetic control of cell size at cell division in yeast. *Nature* 256, 457–451.
- Nurse, P. (1990). Universal control mechanism regulating onset of M-phase. *Nature* 344, 503–508.
- Nurse, P., and Bissett, Y. (1981). Gene required in G1 for commitment to cell cycle and in G2 for control of mitosis in fission yeast. *Nature* 292, 558–560.
- Nurse, P., and Thuriaux, P. (1984). Temperature-sensitive allosuppressor mutants of the fission yeast *Schizosaccharomyces pombe* influence cell-cycle control over mitosis. *Mol. Gen. Genet.* 146, 167–178.
- Ohkura, H., Adachi, Y., Kinoshita, N., Niwa, O., Toda, T., and Yanagida, M. (1988). Cold-sensitive and caffeine-supersensitive mutants of the *Schizosaccharomyces pombe dis* genes implicated in sister chromatid separation during mitosis. *EMBO J.* 7, 1465–1473.
- Ono, Y., Ohno, M., and Shimura, Y. (1994). Identification of a putative RNA helicase (HRH1), a human homolog of yeast Prp22. *Mol. Cell. Biol.* 14, 7611–7620.
- Pines, J., and Hunter, T. (1991). Cyclin-dependent kinases: a new cell cycle motif? *Trends Cell Biol.* 1, 117–121.
- Potashkin, J., and Frendewey, D. (1989). Splicing of the U6 RNA precursor is impaired in fission-yeast pre-mRNA splicing mutants. *Nucleic Acids Res.* 17, 7821–7831.
- Potashkin, J., Li, R., and Frendewey, D. (1989). Pre-mRNA splicing mutants of *Schizosaccharomyces pombe*. *EMBO J.* 8, 551–559.
- Potashkin, J., Naik, K., and Wentz-Hunter, K. (1993). U2AF homolog required for splicing in vivo. *Science* 262, 573–575.
- Reijo, R.A., Cho, D.-S., and Huffaker, T.C. (1993). Deletion of a single-copy tRNA affects microtubule function in *Saccharomyces cerevisiae*. *Genetics* 135, 955–962.
- Rosenberg, G.H., Alahari, S.K., and Käufer, N.F. (1991). *prp4* from *Schizosaccharomyces pombe*, a mutant deficient in pre-mRNA splicing, isolated using genes containing artificial introns. *Mol. Gen. Genet.* 226, 305–309.
- Rush, M.G., Drivas, G., and D'Eustachio, P. (1996). The small nuclear GTPase Ran: how much does it run? *BioEssays* 18, 103–112.
- Russell, P., and Nurse, P. (1986). *cdc25⁺* functions as an inducer in the mitotic control of fission yeast. *Cell* 45, 145–153.
- Russell, P., and Nurse, P. (1987). Negative regulation of mitosis by *wee1⁺*, a gene encoding a protein kinase homolog. *Cell* 49, 559–567.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Schmid, S.R., and Linder, P. (1992). D-E-A-D protein family of putative RNA helicases. *Mol. Microbiol.* 6, 283–291.
- Schwer, B., and Guthrie, C. (1991). PRP16 is an RNA-dependent ATPase that interacts transiently with the spliceosome. *Nature* 349, 494–499.
- Shea, J.E., Toyn, J.H., and Johnston, L.H. (1994). The budding yeast U5 snRNP Prp8 is a highly conserved protein which links RNA splicing with cell-cycle progression. *Nucleic Acids Res.* 22, 5555–5564.
- Spector, D.L. (1993). Macromolecular domains within the cell nucleus. *Annu. Rev. Cell Biol.* 9, 265–315.
- Sugimoto, K., Matsumoto, K., Kornberg, R.G., Reed, S.I., and Wittenberg, C. (1995). Dosage suppressors of the dominant G1 cyclin mutant *CLN3-2*: identification of a yeast gene encoding a putative RNA/ssDNA binding protein. *Mol. Gen. Genet.* 248, 712–718.
- Takahashi, K., Yamada, H., and Yanagida, M. (1994). Fission-yeast minichromosome loss mutants *mis* cause lethal aneuploidy and replication abnormality. *Mol. Biol. Cell* 5, 1145–1158.
- Takeuchi, M., and Yanagida, M. (1993). A mitotic role for a novel fission-yeast protein kinase *dsk1* with cell-cycle stage-dependent phosphorylation and localization. *Mol. Biol. Cell* 4, 247–260.
- Tani, T., and Ohshima, Y. (1989). The gene for the U6 RNA in fission yeast has an intron. *Nature* 337, 87–90.
- Vieira, J., and Messing, J. (1987). Production of single-stranded plasmid DNA. *Methods Enzymol.* 153, 3–11.
- Warbrick, E., and Glover, D. (1994). A *Drosophila* gene encoding a DEAD-box RNA helicase can suppress loss of *wee1/mik1* function in *Schizosaccharomyces pombe*. *Mol. Gen. Genet.* 245, 654–657.
- Wilson, S.M., Datar, K.V., Paddy, M.R., Swedlow, J.R., and Swanson, M.S. (1994). Characterization of nuclear polyadenylated RNA-binding proteins in *Saccharomyces cerevisiae*. *J. Cell Biol.* 127, 1173–1184.
- Wright, A., Maundrell, K., Heyer, W., Beach, D., and Nurse, P. (1986). Vectors for the construction of gene banks and the integration of cloned genes in *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae*. *Plasmid* 15, 156–158.
- Yanez, R.J., Rodriguez, J.M., Bournsnel, M., Rodriguez, J.F., and Vinuela, E. (1993). Two putative African swine fever virus helicases similar to yeast “DEAH” pre-mRNA processing proteins and vaccinia virus ATPases D11L and D6R. *Gene* 134, 161–174.
- Zeitlin, S., Wilson, R.C., and Efstratiadis, A. (1989). Autonomous splicing and complementation of in vivo-assembled spliceosomes. *J. Cell Biol.* 108, 765–777.
- Zeng, C., He, D., Berget, S.M., and Brinkley, B.R. (1994). Nuclear-mitotic apparatus protein: a structural protein interface between the nucleoskeleton and RNA splicing. *Proc. Natl. Acad. Sci. USA* 91, 1505–1509.