

Immunopurification of Ago1 miRNPs selects for a distinct class of microRNA targets

Xin Hong^{a,1}, Molly Hammell^{b,1}, Victor Ambros^{b,2}, and Stephen M. Cohen^{a,2}

^aTemasek Life Sciences Laboratory and Department of Biological Sciences, National University of Singapore, 1 Research Link, Singapore 117604; and ^bProgram in Molecular Medicine, University of Massachusetts Medical School, Worcester, MA 01605

Contributed by Victor Ambros, July 22, 2009 (sent for review June 26, 2009)

microRNAs comprise a few percent of animal genes and have been recognized as important regulators of a diverse range of biological processes. Understanding the biological functions of miRNAs requires effective means to identify their targets. Combined efforts from computational prediction, miRNA over-expression or depletion, and biochemical purification have identified thousands of potential miRNA-target pairs in cells and organisms. Complementarity to the miRNA seed sequence appears to be a common principle in target recognition. Other features, including miRNA-target duplex stability, binding site accessibility, and local UTR structure might affect target recognition. Yet computational approaches using such contextual features have yielded largely nonoverlapping results and experimental assessment of their impact has been limited. Here, we compare two large sets of miRNA targets: targets identified using an improved Ago1 immunopurification method and targets identified among transcripts up-regulated after Ago1 depletion. We found surprisingly limited overlap between these sets. The two sets showed enrichment for target sites with different molecular, structural and functional properties. Intriguingly, we found a strong correlation between UTR length and other contextual features that distinguish the two groups. This finding was extended to all predicted microRNA targets. Distinct repression mechanisms could have evolved to regulate targets with different contextual features. This study reveals a complex relationship among different features in miRNA-target recognition and poses a new challenge for computational prediction.

Argonaute | gene regulation | RISC complex

Animal genomes contain hundreds of microRNA genes (miRBase 13.0). Recent estimates suggest that miRNAs comprise $\approx 1\%$ of genes in *Drosophila* and *Caenorhabditis elegans* and 2–3% of genes in mouse and human. To date, functional analysis in vivo has revealed biological roles for only a small fraction of these (1–3). One issue limiting progress in understanding the miRNA functions is identification of the target mRNAs that they regulate. Computational target identification is primarily based on sequence complementarity to the miRNA (reviewed in ref. 2), but many computational strategies also make use of sequence context to predict miRNA targets. Comparisons of different methods show limited overlap among the predicted targets, although those that place more emphasis on pairing to the seed sequence at the 5' end of the miRNA tend to produce similar results. Most of these methods identify many possible targets for each miRNA, often hundreds (e.g., refs. 4–9).

A growing body of experimental evidence shows that miRNAs can regulate many targets. Overexpression of miRNAs in heterologous cell types can affect the levels of hundreds of mRNAs with target sites (e.g., ref. 10). Conversely, depletion of miRNAs can lead to increased levels of a comparable number of target mRNAs (11, 12). Changes in target RNA stability can result from miRNA-induced deadenylation of the mRNA (13–15). mRNA up-regulation, combined with target prediction has helped to identify biologically relevant targets of specific

miRNAs (e.g., ref. 16). Whole proteome analyses have shown that miRNA induced changes in protein expression correlate with changes in mRNA level, in trend if not in magnitude (17, 18). Yet, there are well-documented instances of miRNA-mediated regulation at the protein level that do not involve changes in mRNA level (14, 17, 18). Therefore, methods to identify targets by miRNA-induced changes in expression profile can only tell part of the story. This highlights the need for alternative means to identify miRNA targets.

One such alternative involves identification of microRNA targets by virtue of their physical association with miRNA-containing ribonucleoprotein complexes (19–24). In ref. 19, we reported a method based on Ago1 immunopurification (IP) that proved to be effective. Eleven new targets were identified for miR-1, including some that had not been predicted. Although the specificity was high, with all new targets experimentally validated, the method had limited sensitivity, identifying $\approx 1/10$ th of the expected number of targets. Here, we present an improved Ago1 IP protocol, which permits identification of hundreds of potential miRNA targets, and compare the contextual features of targets identified by IP to the targets destabilized at the mRNA level upon Ago1 depletion.

Results

In an effort to improve the sensitivity of miRNA IP, with minimal loss of specificity, we tested a variety of antibody concentrations, incubation times and wash conditions (Fig. S1). Sensitivity was assessed by quantitative PCR to monitor miRNA levels (over a broad range of abundance: miR-184 comprises 17% of S2 cell miRNA; miR-305: 1.5%; miR-7: 0.1%; miR-92b: 0.01%; (25)). The 4 miRNAs were enriched >50 -fold in IP from cells expressing HA-tagged Ago1 compared with control cells not expressing the transgene (Fig. 1A, *, $P < 0.05$). The small nucleolar RNA (snoR227) showed no enrichment ($\log_2 = 0.18$). Thus, the IP protocol can recover miRNAs over a broad range of expression levels without enriching for unrelated small RNAs.

The association between miRNAs and targets must be stable for IP to be useful. To assess this association, we measured the recovery of *reaper* mRNA, a known target of miR-2a. The miR-2 family comprises $>13\%$ of S2 cell miRNAs. *reaper* mRNA was enriched ≈ 32 -fold by IP. A control lacking known miRNA binding sites was not enriched (Fig. 1B). This compares favorably with the 2- to 3-fold enrichment of *reaper* obtained in the study in ref. 19.

Author contributions: X.H., M.H., and S.M.C. designed research; X.H. and M.H. performed research; X.H., M.H., V.A., and S.M.C. analyzed data; and X.H., M.H., V.A., and S.M.C. wrote the paper.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

¹X.H. and M.C.H. contributed equally to this work.

²To whom correspondence may be addressed. E-mail: cohen@tll.org.sg or vrambros@gmail.com.

This article contains supporting information online at www.pnas.org/cgi/content/full/0908149106/DCSupplemental.




Fig. 1. miRNP immunopurification. (A) Enrichment of miRNAs measured by Q-PCR. y axis: fold enrichment (log₂ scale). Splicing RNA *U27* was used for normalization and *snoR227* as a control. Results represent 3 independent IP experiments. (B) Enrichment of a known miR-2a target *reaper* by IP. Data normalized to *rp49* for 3 independent experiments (*t* test, $P < 0.05$).

Expression Profiling of mRNA Associated with Ago1. To examine the population of mRNAs associated with Ago1 complexes, total RNA was recovered by IP with anti-HA from S2 cells expressing HA-Ago1 and from untransfected control cells. After 2 rounds of linear amplification the RNA was used to generate probes for expression profiling. Approximately 7,000 transcripts can be detected reliably by expression profiling in S2 cells (11). A total of 1,621 mRNAs were significantly IP enriched with a P value of < 0.05 . Of these, 1,191 were enriched > 1.4 -fold, 464 were enriched > 2 -fold and 89 were enriched > 4 -fold (Fig. S2A). A considerable fraction of S2 cell transcripts are associated with Ago1 complexes, although for most the degree of enrichment is low in magnitude.

S2 cells miRNAs fall into 26 seed families with distinct target specificities (Table S1). The 5 and 10 most abundant seed families comprise 73% and 88% of S2 cell miRNA. IP-enriched targets for all S2 miRNAs were slightly more abundant than S2 cell transcripts in general ($P < 1 \times 10^{-10}$; Table S2 and Fig. S2B). IP-enriched targets for the top 5 and top 10 seed families were also slightly more abundant ($P = 0.02$; $P = 3.5 \times 10^{-6}$). Although statistically significant, these differences are small in magnitude and should not pose a concern for use of IP for target identification.

We next compared IP using the Ago1 transgene with IP of endogenous Ago1 using monoclonal anti-Ago1 [kindly provided by M. Siomi (Keio University School of Medicine, Tokyo, Japan)]. Enrichment was determined relative to a control IP with monoclonal anti-Myc after normalization to *rp49*. Sixteen targets were assayed by quantitative RT-PCR (Q-PCR). All showed significant enrichment, with a strong correlation between the two methods (Fig. S1C, correlation coefficient $r = 0.71$, $P = 0.0018$). Thus, the results are largely independent of the antibody used.

Experimental Validation of Target Enrichment. The performance of Ago1 IP was tested with reference to a nonredundant set of experimentally validated miRNA-target pairs, consisting of 67

validated positives and 29 validated negatives (Table S3). More targets were IP-enriched than expected after normalization for the number of transcripts in each category (Fig. 2A). Enrichment was > 3 -fold for targets of the 5 and 10 most abundant miRNA seed families (20 positives/489 IP-enriched with sites for the top 10 miRNAs; $P = 6.0 \times 10^{-5}$). The enrichment was slightly lower when all S2 cell miRNAs were considered (23 positives/745 IP-enriched transcripts with sites for all miRNAs), but remained significant ($P = 3.6 \times 10^{-4}$). There was no enrichment for the 29 miRNA-target pairs that tested negative experimentally. For comparison with the analysis of *reaper* (Fig. 1B), the average magnitude of IP enrichment was 5.5-fold for all validated positives. Enrichment was greatest for abundant miRNAs (Fig. S1D). It is noteworthy that some validated targets were not enriched by IP. Different factors may contribute to this. Some in vivo validated pairs might not be functional in S2 cells. Among the validated targets for top 5 miRNAs that were not IP enriched, 28% were up-regulated at the RNA level upon Ago1 depletion (Table S3). Target turnover may also affect recovery.

Antisense oligonucleotides were used to deplete specific miRNAs in S2 cells to ask if target recovery depends on the miRNA. miR-184 was chosen because many of its predicted targets were enriched by IP (Table S4). Cells were treated with anti-miR-184 or with a scrambled sequence control and subjected to IP with anti-Ago1. Nine of the thirty-two miR-184 targets were assayed by Q-PCR. IP recovery of 7 was lower in miR-184 depleted cells (Fig. 2B), indicating that binding to miR-184 contributes to IP enrichment. The partial reduction in recovery likely reflects incomplete depletion of miR-184. Other miRNAs might also contribute: 8/9 have predicted sites for miRNAs other than miR-184, including sites for 4 of the top 10 seed families. IP enrichment of specific targets can be attributed to association with a specific miRNA.

Additional Validation of New miRNA Targets Identified by IP. From the set of computationally predicted miR-184 targets, we selected 17 that were IP enriched and 17 that were not. Transcript levels were measured by Q-PCR after miR-184 depletion. Twelve of seventeen IP enriched targets were up-regulated in cells depleted of miR-184 ($P < 0.05$; Fig. S3A and Table 1). Control mRNAs were not affected. For comparison, only 2 of a set of 17 miR-184 targets that were not IP-enriched were up-regulated in miR-184 depleted cells. Luciferase reporter assays were also used to assess miRNA-mediated regulation of the set of IP enriched miR-184 targets. Five of the twelve targets up-regulated on miR-184 depletion have been validated (7). We tested one of these as a positive control, 5 other up-regulated transcripts and 3 from the IP, but not up-regulated set (Table 1). Control UTRs lacking miR-184 sites were unaffected by miR-184 depletion. The 6 up-regulated transcripts showed modest




Fig. 2. Transcripts associated with Ago1 complexes. (A) Comparison of IP results with experimentally validated miRNA targets. Fold enrichment of the nonredundant target set (Table S3) for the 5 and 10 most abundant miRNA families and for all S2 miRNA families. y axis: enrichment calculated as the number of validated targets IP-enriched divided by the number not IP-enriched after normalization for transcript number in each category. P values: Fisher's exact two-tailed test. (B) Effect of miR-184 depletion on the recovery of predicted targets. y axis: Relative IP enrichment calculated by normalizing IP fold enrichment in miR-184 depleted cells to control cells. Transcript levels quantified by Q-PCR (normalized to *rp49*) and compared with control IP with empty beads. *reaper* lacks miR-184 sites and serves as a control. *, $P < 0.05$, Student's *t* test.




Fig. 4. A comparative analysis on Ago1 IP-enriched transcripts Vs Ago1 RNAi-up-regulated transcripts. (A) Numbers of mRNAs with target sites identified by IP or by Ago1 RNAi. IP-enriched used a cutoff >1.4 -fold and $P < 0.05$. Ago1 RNAi set from (12): up-regulated >1.5 -fold ($P < 0.05$). All: all S2 cell mRNAs; all S2 miRNA: those with sites for any S2 miRNA. (B) y axis: median UTR length (nt) represented by gray squares. The range between 1st and 3rd quartiles are indicated by bars. P values: Wilcoxon U test, Bonferroni corrected (4 tests per group). The median for the Ago1 RNAi set is 262 nt longer than the IP set ($P = 6e-21$) and 187 nt longer than for all RNAs with sites ($P = 6e-13$). The IP set was 75 nt shorter than all RNAs with sites ($P = 5 \times 10^{-7}$). (C) Site density profiles in IP-enriched vs. Ago1 RNAi-up-regulated targets and all S2 miRNAs. The up-regulated set has more sites/transcript than the IP set (median shift 4, $P = 2 \times 10^{-18}$) or than all RNAs with sites (median shift 3, $P = 1 \times 10^{-14}$). Median site density was 6.9/500 nt in the up-regulated set vs. 6.0 for the IP set and 5.6 for all RNAs. These differences were significant using a K-S test (two tail). (D) Assessment of site openness for the Ago1 RNAi up-regulated set vs. all RNAs with sites (as in Fig. 3B). (E) Fold enrichment for the optimal upstream and downstream windows in IP-enriched and Ago1 RNAi up-regulated sets. x axis: average nucleotide openness binned from 10% to 80%. y axis: fold enrichment as compared with all S2 transcripts with sites, which were used as the common control. Pearson's correlation coefficient, r values and associated P values are shown.

energy of miRNA-target base pairing for transcripts enriched by IP with those not enriched (ΔG_{hybrid} as in ref. 9). The distribution of binding energies was significantly shifted toward more stable duplexes in IP-enriched transcripts ($P = 0.005$ using a two-sample Kolmogorov–Smirnov test). The correlation between binding energy ΔG_{hybrid} and degree of enrichment in the IP set was modest ($r = -0.60$), on the borderline of statistical significance ($P = 0.0503$). On this basis, IP does not appear to select strongly for more stable target site binding.

Local secondary structure in the UTR might affect function by making a miRNA site more or less accessible. Previous studies have developed ways to predict site “openness” and suggested that this could improve computational target prediction (7, 9, 33). We found a significant shift toward greater site openness (9) in IP-enriched targets compared with those not enriched (Kolmogorov–Smirnov two-sample, $P = 4 \times 10^{-6}$; Table S6). The degree of enrichment for openness correlated well with increasing site accessibility ($r = 0.865$, $P = 0.006$). A comparable correlation was found when all S2 mRNAs were used as the control set ($r = 0.868$, $P = 0.005$).

To examine the impact of nearby sequences, we tested windows from 5 to 95 nt in length upstream and downstream of the site (Table S7). Fig. 3B presents a surface plot showing the correlation coefficient between degree of openness and enrichment in the IP set compared with all S2 transcripts. Upstream of the site, the correlations were significant for windows of up to 50 nt. Downstream, the correlations were significant for windows from 10 to 70 nt with an optimum at 50 nt (Table S7 and Fig. 3B).

A previous analysis using a different miRNP protein in *C. elegans* (AIN-IP, GW182) found that a 25-nt upstream window had the best correlation with IP enrichment, but did not find evidence for downstream openness (9). These studies may have sampled different miRNP complexes, since antibodies to different RISC components were used, but species differences in UTR composition cannot be ruled out.

Comparison of Targets Identified by Ago1 IP and Ago1 Depletion.

Expression profiling of cells depleted of miRNA function is gaining acceptance as a means of identifying potential targets (e.g., refs. 11, 12, 16, and 18). In this context, we asked whether similar target features are selected for by IP and Ago1 depletion. A priori we expect them to be similar, because both reflect activity of Ago1 complexes. To our surprise, we found $<1/3$ overlap in the RNAs identified by the two methods (Fig. 4A and Table S8 lists all transcript data). Recent reports using Ago-2 to pull down miRNA targets in mammalian cells over-expressing a miRNA also showed limited overlap with targets downregulated at the mRNA level (20, 23). To explore the basis for these differences we examined the two groups for distinct structural and molecular features, using all S2 mRNAs as a common control set.

There was little difference in seed types enriched by the two methods (Fig. S4A). However, sites with stronger binding energy were under-represented in the up-regulated transcript set ($P = 0.01$, cumulative distribution plot in Fig. S4B). There was a significant anti-correlation between more stable hybrids and

proaches that favor site context, overall binding energy or conservation are likely to favor one part of the UTR target spectrum over another. A systematic analysis of the interactions between contextual features, including synergistic regulation and structural openness, could improve the accuracy of prediction.

Materials and Methods

Immunopurification of *Ago1* from S2 cells is described in Fig. S1A and *SI Materials and Methods*. UTR luciferase reporter assays were performed as described in refs. 27 and 38 with modifications described in *SI Materials and Methods*. miRNA and mRNA quantitative PCR was performed as described in *SI Materials and Methods*. Expression profiling was performed by the EMBL

Gene Core using Affymetrix *Drosophila* 2.0 arrays (see *SI Materials and Methods*). miRNA target site prediction and statistical analysis is detailed in *SI Materials and Methods*. David bioinformatic tools were used for gene ontology. Functional annotation clustering was applied as described in ref. 39.

ACKNOWLEDGMENTS. We thank V. Benes and T. Bähr-Ivacevic of the EMBL GeneCore for expression profiling; S. Lim and E. Loeser for technical support; T. Sandmann for help with microarray and bioinformatic analysis; and J. Varghese, W. Ge, P. Verma for helpful suggestions. Work in the S.M.C. laboratory was supported by European Union Sixth Framework Programme Grant "Sirocco" LSHG-CT-2006-037900 and the Singapore Millennium Foundation via core funding to Temasek Life Sciences Laboratory. Work in the V.A. laboratory was supported by National Institutes of Health Grants F32GM087039 (to M.H.) and GM34028 and GM066826 (to V.A.).

1. Bushati N, Cohen SM (2007) microRNA functions. *Annu Rev Cell Dev Biol* 23:175–205.
2. Bartel DP (2009) MicroRNAs: Target recognition and regulatory functions. *Cell* 136:215–233.
3. Miska EA, et al. (2007) Most *Caenorhabditis elegans* microRNAs are individually not essential for development or viability. *PLoS Genet* 3:e215.
4. Stark A, Brennecke J, Bushati N, Russell RB, Cohen SM (2005) Animal MicroRNAs confer robustness to gene expression and have a significant impact on 3' UTR evolution. *Cell* 123:1133–1146.
5. Lewis BP, Burge CB, Bartel DP (2005) Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120:15–20.
6. Krek A, et al. (2005) Combinatorial microRNA target predictions. *Nat Genet* 37:495–500.
7. Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E (2007) The role of site accessibility in microRNA target recognition. *Nat Genet* 39:1278–1284.
8. Kheradpour P, Stark A, Roy S, Kellis M (2007) Reliable prediction of regulator targets using 12 *Drosophila* genomes. *Genome Res* 17:1919–1931.
9. Hammell M, et al. (2008) mirWIP: microRNA target prediction based on microRNA-containing ribonucleoprotein-enriched transcripts. *Nat Methods* 5:813–819.
10. Lim LP, et al. (2005) Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* 433:769–773.
11. Rehwinkel J, et al. (2006) Genome-wide analysis of mRNAs regulated by Drosha and Argonaute proteins in *Drosophila melanogaster*. *Mol Cell Biol* 26:2965–2975.
12. Eulalio A, et al. (2007) Target-specific requirements for enhancers of decapping in miRNA-mediated gene silencing. *Genes Dev* 21:2558–2570.
13. Giraldez AJ, et al. (2006) Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. *Science* 312:75–79.
14. Behm-Ansmant I, et al. (2006) mRNA degradation by miRNAs and GW182 requires both CCR4:NOT deadenylase and DCP1:DCP2 decapping complexes. *Genes Dev* 20:1885–1898.
15. Eulalio A, et al. (2009) Deadenylation is a widespread effect of miRNA regulation. *Rna* 15:21–32.
16. Karres JS, Hilgers V, Carrera I, Treisman J, Cohen SM (2007) The conserved microRNA miR-8 tunes atrophin levels to prevent neurodegeneration in *Drosophila*. *Cell* 131:136–145.
17. Selbach M, et al. (2008) Widespread changes in protein synthesis induced by microRNAs. *Nature* 455:58–63.
18. Baek D, et al. (2008) The impact of microRNAs on protein output. *Nature* 455:64–71.
19. Easow G, Teleman AA, Cohen SM (2007) Isolation of microRNA targets by miRNP immunopurification. *Rna* 13:1198–1204.
20. Karginov FV, et al. (2007) A biochemical approach to identifying microRNA targets. *Proc Natl Acad Sci USA* 104:19291–19296.
21. Zhang L, et al. (2007) Systematic identification of *C. elegans* miRISC proteins, miRNAs, and mRNA targets by their interactions with GW182 proteins AIN-1 and AIN-2. *Mol Cell* 28:598–613.
22. Ender C, et al. (2008) A human snoRNA with microRNA-like functions. *Mol Cell* 32:519–528.
23. Hendrickson DG, Hogan DJ, Herschlag D, Ferrell JE, Brown PO (2008) Systematic identification of mRNAs recruited to argonaute 2 by specific microRNAs and corresponding changes in transcript abundance. *PLoS ONE* 3:e2126.
24. Landthaler M, et al. (2008) Molecular characterization of human Argonaute-containing ribonucleoprotein complexes and their bound target mRNAs. *Rna* 14:2580–2596.
25. Ruby JG, et al. (2007) Evolution, biogenesis, expression, and target predictions of a substantially expanded set of *Drosophila* microRNAs. *Genome Res* 17:1850–1864.
26. Doench JG, Sharp PA (2004) Specificity of microRNA target selection in translational repression. *Genes Dev* 18:504–511.
27. Brennecke J, Stark A, Russell RB, Cohen SM (2005) Principles of microRNA-target recognition. *PLoS Biol* 3:e85.
28. Varghese J, Cohen SM (2007) microRNA miR-14 acts to modulate a positive autoregulatory loop controlling steroid hormone signaling in *Drosophila*. *Genes Dev* 21:2277–2282.
29. Nahvi A, Shoemaker CJ, Green R (2009) An expanded seed sequence definition accounts for full regulation of the *hid* 3' UTR by bantam miRNA. *Rna* 15:814–822.
30. Simon DJ, et al. (2008) The microRNA miR-1 regulates a MEF-2-dependent retrograde signal at neuromuscular junctions. *Cell* 133:903–915.
31. Johnson SM, et al. (2005) RAS is regulated by the let-7 microRNA family. *Cell* 120:635–647.
32. Vella MC, Choi EY, Lin SY, Reinert K, Slack FJ (2004) The *C. elegans* microRNA let-7 binds to imperfect let-7 complementary sites from the *lin-41* 3' UTR. *Genes Dev* 18:132–137.
33. Long D, et al. (2007) Potent effect of target structure on microRNA function. *Nat Struct Mol Biol* 14:287–294.
34. Grimson A, et al. (2007) MicroRNA targeting specificity in mammals: Determinants beyond seed pairing. *Mol Cell* 27:91–105.
35. Cairrao F, Halees AS, Khabar KS, Morello D, Vanzo N (2009) AU-rich elements regulate *Drosophila* gene expression. *Mol Cell Biol* 29:2636–2643.
36. von Roretz C, Gallouzi IE (2008) Decoding ARE-mediated decay: Is microRNA part of the equation? *J Cell Biol* 181:189–194.
37. Trabucchi M, et al. (2009) The RNA-binding protein KSRP promotes the biogenesis of a subset of microRNAs. *Nature* 459:1010–1014.
38. Bushati N, Stark A, Brennecke J, Cohen SM (2008) Temporal reciprocity of miRNAs and their targets during the maternal-to-zygotic transition in *Drosophila*. *Curr Biol* 18:501–506.
39. Huang DW, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4:44–57.