

# A sequence compilation and comparison of exons that are alternatively spliced in neurons

Stefan Stamm\*, Mike Q.Zhang, Thomas G.Marr and David M.Helfman

Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA

Received January 28, 1994; Accepted March 16, 1994

## ABSTRACT

**Alternative splicing is an important regulatory mechanism to create protein diversity. In order to elucidate possible regulatory elements common to neuron specific exons, we created and statistically analysed a database of exons that are alternatively spliced in neurons. The splice site comparison of alternatively and constitutively spliced exons reveals that some, but not all alternatively spliced exons have splice sites deviating from the consensus sequence, implying diverse patterns of regulation. The deviation from the consensus is most evident at the –3 position of the 3' splice site and the +4 and –3 position of the 5' splice site. The nucleotide composition of alternatively and constitutively spliced exons is different, with alternatively spliced exons being more AU rich. We performed overlapping k-tuple analysis to identify common motifs. We found that alternatively and constitutively spliced exons differ in the frequency of several trinucleotides that cannot be explained by the amino acid composition and may be important for splicing regulation.**

## INTRODUCTION

Among all animal tissues the brain is probably the most molecularly complex organ with about 30% of the mammalian genome expression dedicated to it [1]. Both kinetic [2] and clonal [3] analysis indicate that 40–65% of the mRNA expressed in brain is restricted to this tissue. Furthermore, the majority of this mRNA is expressed in neurons and not in glia [4], [5]. Neuron specific gene expression can be achieved by transcriptional (reviewed in [5, 6]) and post transcriptional mechanisms including splicing and RNA editing [7]. In order to analyze common features of neuron specific exons, we compiled and analyzed the currently available neuron specific exons.

It has been estimated that alternative splicing is involved in more than 5% of on/off regulation in *Drosophila* genes [8]. In addition, it has been shown that the inclusion of alternatively spliced exons alters the electrophysiological properties of ion channels like the Glutamine A–D [9] and the NMDA [10] receptors. In 11 genes listed in this survey, alternatively spliced

exons encode stop codons leading to truncated proteins. In 10 genes listed, the alternatively spliced exon introduces a phosphorylation site and in one case a phosphorylation site is removed.

The exact mechanisms that regulate the alternative use of neuron specific exons are not well understood. Several regulatory sequences have been identified, including splice sites deviating from the consensus [11], RNA binding factors [12, 13], elements in the flanking introns [14, 15] and secondary structures [16]. In addition, several components of the splicing machinery have been found to be specific for neurons, among them the composition of SR proteins [17], U2 RNA composition [18] and the protein SmN [19] that might be involved in the etiology of the Prader Willi syndrome [20], but whose functional significance is not clear [21].

Neuron specific alternatively spliced exons can be the result of neuron specific transcription followed by alternative splicing, or the result of transcription that takes place in all cells followed by neuron specific splicing. Although the consequence in both cases is an exon that is alternatively spliced in neurons, the mechanistic regulation might be quite different. For example, neuron specific exons generated by neuron specific promoter use are alternatively regulated in non-neuronal cells [22]. In contrast, several genes exhibit neuron specific splicing only when expressed in neuronal cells, but not in any other cell type [11], [15, 23]. The neuron-specific usage of the exons compiled here has been established by comparison with some, but not all, non-neuronal tissues like liver or muscle. Therefore, more detailed investigations may detect some use of these exons in non-neuronal cell types. Furthermore, inspection of the compiled sequences shows that certain functional subclasses such as receptors are highly represented, which probably reflects the current focus of research on these molecules, rather than their greater use of alternative spliced exons.

## METHODS

### Collection of alternative spliced exons and control database

Exons that are alternatively spliced in neurons were collected from the literature using the Medline database, searching with the key words 'alternative splicing' and 'brain' or 'neuron'. The sequences were run against GenBank using the BLASTN 1.3.12

\*To whom correspondence should be addressed

**Table 1.** Compilation of exons that are alternatively spliced in neurons

Gene	regulation	experimental evidence	accession number	species	sequence	ref.
Glycine receptor $\alpha 2A$	regional and developmental	neuronspecific, in situ	X57281	R	ttgcattctcgag#GTCCCTCAGTAAACGTTACTTGCAATATTATCACACAGTTGGATCG TCACAGAACACCACATGgtaaatgtcactagt	[50]
Glycine receptor $\alpha 2B$	regional and developmental	neuronspecific, in situ	X61159	R	gttcaattacatgg#GGCCCTCTTAAATGGTACCTGCAACATATTGGGTAATAGCAGAAA CAATGGtgatgtggactgatccatgt	[50]
Glutamate receptor, B type, flop	regional and developmental	neuronspecific, in situ, electrophysiology	M36419	R	atgttttatcggttaaga@ATGGGGTTAACCTCGCAGTACTAAACTGAATGAAACAAGGCTG TTGGGAAATTGAAAACAAATGGTGGTACGACAAGGAGTCGGCAGCGGGGAGGTGATT CAAGtgtacgcccc	[9]
Glutamate receptor, B type, flip	regional and developmental	neuronspecific, in situ, electrophysiology	M38061	R	tattttccacgtgag@ACCCCGAGTAAATCTTGAGTATTGAAACTCAGTGAGCAAGGGCTT AACGACAAGCTGAAAACAAATGGTGGTACGATAAAGGTGAATGTGGAGCCAAGGACTCGGGAAAG TAAGtgtacgttcgtcgag	[51]
rat brain Ca channel, alpha subunit rbc-1		expression in brain, heart and adreanal gland	M67516	R	#GGTTACTTTAGTGATCCCTGGATGTTTGACTTCCTCATCGTCAATTGGGAGCATATAATGATG TCATTCCTCAGTAACTAATgtgatgt	[52]
rat brain Ca channel, alpha subunit rbc-1			M89924	R	ccccatgcg@CACTATTCTGTGATGCCATGGAAATACATTGACGCCCTGATTGTTGGTAGC ATTGTTGATATAGCAATCACCAGGGTACAC	[52]
Ca channel $\alpha 1$ , form 1		expression in brain and cardiac muscle, heart Northern analysis	L01776	M	#ATGCAAGAGCTATGCCATGGATGAGTGCCTGGGTGATTTGTCAGTCGGTACATCTGGAT CCCTTTCTGCTCTAAATCTGGTTCTGGGTTG	[53]
Ca channel $\alpha 1$ , form 1		expression in brain and cardiac muscle, heart Northern analysis	L01776	M	#GTCATGATGCCGAGGAAAGGACTGGCCCTGGATCTATTGTAACACTAACATCATAGGG CATTTTTGACTTACTTGTGTTCTGGGTTGAGCGGGAGTTTCAAGAGGAGGGAGAAA GTCAAAGGCCGGAGGAGATTCCAGAGCTTCAGAGGAGAACGACACTAGAGAAAGATCTCAAAGG CTTACCTGAGATGCCACCCAGGCAAGAACATTGACCCCCGAGAATGAGGAGCGAGGGCATGGATG AAGACAAAGCTCGAACACA	[53]
Ca channel $\alpha 1$ , form 4		expression in brain and cardiac muscle, heart Northern analysis	L01776	M	#GGTTACTTTAGTGATCCCTGGATGTTTGACTTCCTCATCGTCAATTGGGAGCATATAATGATG TCATTCCTCAGTAACTAATgtgatgt	[53]
		expression in brain and cardiac muscle, Northern analysis	L01776	M	#CACTATTCTGTGATGCCATGGAAATACATTGACGCCCTGATTGTTGGTAGCATTGATA TAGCAATCACCAGGGTACAC	[53]
R15 cDNA	regional (ring ganglion)	neuron specific, in situ, protection	M17535	A	#TATATG	[54]
	regional (abdominal ganglion)	neuron specific, in situ, protection	M17536	A	#AGTGATGGAAGCGCAGAGAGAACCGTACACCAAGGATGGGATCCGGG	[54]

cassette exons, expression of the gene in all tissues, expression of the exons only in neurons

Clathrin light chain B	developmental	expression in neurons, PCR, immunocytochemistry	L01564	R	tcctttctcaacggtttcccaagG@ATCGCTGACAAAGCGTTCTACCAGCAGGCCAGATGCTGA TACCATGGCTATGTgtaatcggtcccttgc	[51] [55]
Clathrin light chain B		expression in neurons, immunocytochemistry	M20469	H	G@ATCGCTGACAAAGCATTCTACCAGCAGGCCAGATGCTGATATCATCGGCTACGT	[56]
Clathrin light chain B			X04852	B	G@ATCGCTGACAAAGCATTCTACCAGCAGGCCAGATGCTGATATCATCGGCTACGT	[56]
Clathrin light chain B			N/A	M	gctgttagG@ATCGCTGACAAAGCGTTCTACCAGCAGGCCAGATGCTGATACCATTGGCTATGTg tatgtgca	[57]
Clathrin light chain A, LCA2		expression in neurons, Western blot, immunocytochemistry	M20471	R	G@GTGGCAGATGAAGCTTCTACAAAACAACCCCTCGCTGACGTGATTGGTTATGT	[55] [58]
Clathrin light chain A, LCA2			M20471	H	G@GTGGCAGATGAAGCTTCTACAAAACAACCCCTCGCTGACGTGATTGGTTATGT	[56]
Clathrin light chain A, LCA2			X04849	B	G@GTGGCAGATGAAGCTTCTACAAAACAACCCCTCGCTGACGTGATTGGTTATGT	[59]
Clathrin light chain A, LCA1		neuronspecific, immunocytochemistry	M15882	R	C@AACATAAAACCATCCTTGCTACAGCCTAGAACAGG	[55] [58]
Clathrin light chain A, LCA1			M20471	H	C@AACATAAAACCATCCTTGCTACAGCCTAGAACAGG	[56]
Clathrin light chain A, LCA1			X04849	B	C@AACATAAAACCATCCTTGCTACAGCCTAGAACAGG	[59]
neuronal cell adhesion molecule, exon 18	developmental	neuron specific immunocytochemistry	M15939	C	tctttttcacagG@CACACCGCGTACTGCAGCTACTGTTGAGGACATGCTGCCCTCTGTAAC ACGGCACCCACCAACTAATCTGACACTATCTACAGTAAACTTTTGCACTGCTCAGAACAGCCCCACGAG CGAACCCACCCCCACCTGACCTTCAGTATTGCCCAGCAGCCACGCCATACCTGACTCAAAACGCCA TGTCGCTTGCCAGGCTACTCCAGCAGCCAAAGGGGAGGCTCCCCCTGCTCTCCACCAACCCCTCC TCTACGCCAAAGGTGCCCCCTTGTGATCTCAGGATACCCCAAGCTCTGCTCAGCTACTAA TAATTGTCTTCAAGTGTCTGTGCAACCAAAGGTGCACTGCTGAGGCCAGACTGTGCTAAAC TGGCCAGACACTTCAAGCAGCAGCTGGTCACTGAGCTCAGCTGCTCCACCAACCCCTGCAAAACCTCACT AGTCTCAGCTCAGCTCAGGCCAACAGCAGGAGGTCTCAAGCACCAAGAGCCCGAGGAAAGAAC TGGCCAGCCCCAACAGAGCTGAAGAGGCCAACAGAGCCAAAGTCCAAGGCACTGGAGTG AGGCCTGCTTCAGCGCGCACCAACCAACCCCTCCAGAATGGAGCTTAAATGGCACAGGGACC TTCAAGACCCAGACATTTGATCTTGTGAAAGATGTTTTCAGCTTGGCACTACTACTCTG CAGTGTGGCTAGTGGCAAGCTCGTGGCTGAGCTTCTCCACTGAGACAGCTCTGACCTCTG CACCTGCAAAAGCCAGatatggctgcct	[60]
neuronal cell adhesion molecule, VASE exon	developmental	neurons specific, also in heart and adrenal gland	M32611	R	tttgtttctccag@GCATCGTGGACTCGACCAAGAGAAGCAAGAGgtatagtttacctgaccacta gcca	[61], [62]
neuronal cell adhesion molecule, VASE (π)exon	.		X14527	M	ttcttctccag@GCATCGTGGACTCGACCAAGAGAAGCAAGAGgtata	[63]
protein 4.1, exon 15		predominantly expressed in brain, PCR analysis	L00919	M	ttgaatttggcaacgcg@AACCTTGCAGAAAAAGGTGAAGATCTGATAAGAAATGAGGAAGgtt agccatttttcccttc	[64]
Protein 4.1, exon 5		predominantly expressed in brain, PCR analysis	L00919	M	tgttttccacag@AAACATGCTAATTCTAACAGACTTGTGAGGAGGTGCGAGCACCTCAACCT TTTGAGAAGAACGACTACTTGTGTTGACCTTGAGGACAGCCAACTCTAAAGgttaaggagac	[64]
n1 src		neuron specific, Western analysis, PCR	M61224	C	tgcgttcatgttagG@AGAAAGTGGACGTGAGgtgttacccgg	[55], [65]
n1 src		neuron specific, PCR	N/A	F	G@ACGAAGATAACTCGAC	[66]

(Table 1. *Cont.*)



cassette exons, expression of the gene in neuronal tissue where it is alternatively spliced

(Table 1. *Cont.*)

**alternative 3' splice site**  all genes are only expressed in neuronal tissue, the alternatively spliced exon is underlined

GlyReceptor α1	regional developmental	neuron specific immunocytochemistry		R	cttatttttaagtag#AGCCCCATGCTAAATCTGTTTCAGGATGATGAGGGTGG	[110]
NMDA Receptor 1, exon 1		neuron specific, electrophysiology	X63255	R	CTGAGACGCCCGCCGCCCTCTGCCCTCCCCGCAGACAGACGCCACGGGACAGGGCGCTG GCCCAAGCAGACGCCCGACGCCACGGGCTGGGGAGGGACC	[110]
D3 receptor, D3S	regional	neuron specific pharmacological profile	X67274	M	tccccccatcg#AGCCATATAGAACACAAACCATATTCCTCCAGAAATGCCAGGACCCCTCTGTC TCACATCTACAGCCCTCTCTCTGGCAGACAT	[111]

(Table 1. *Cont.*)

D3 receptor(o2del)		neuron specific, electrophysiology	M69192	R	accacagGGGAGATCCCAGGATCTGCCTCCATCTCCAACCCGTATTGTCATTTACTCTTCAGTGG TGTCCCTCTACGTTCCCTCGGGTGACTGTCTGGTC	[84]
POMPC		neuron specific	J00291	H	cattgttttgcccttgcaggGGTCCCCACGAACTCTAGTTGCTCTGCA	[112]
prolactin	regional	expressed in pituitary neurons immunocytochemistry	J00767	R	cactgtacacctgaatttcttagCAG	[113]
nervous system specific RNP protein-1	developmental	neuron specific, in situ	M34894	X	attactttataaaacaagGCGTCCACTCCAACCCGCACAGGGGGTTCTTGGAAC	[114]
		exon introduces frameshift				
Synapsin I	regional	neuron specific, immunocytochemistry	M27812	R	ctttgtctctctcttagC <del>AAA</del> ATCCCAGCTCTGACCAATGCCCTCAACCTTCCAGAGCCAGCCCC	[115]
Synapsin I	regional		J05431	H	tccacaccttgcctctcttagC <del>AAA</del> ATCCCAGCTCTGACCAATGCCCTCAACCTTCCAGAGCCAGCC	[116]



### alternative 5'ends

$\gamma$ -aminobutyric acid A receptor $\beta 4'$ subunit raw2 (NGK2, Kv3.1 b)		neuron specific, electrophysiology	X56647 Y07521	C M	GGTAGAGAGCAGGttttccccc GTGGAGAAACCTCTCAGGGCATGTCGATCTGACCTTTCACCTCCGCCCCCTGTAGCAATGATT CAGATCCACTCAGACTTGCTT	[117] [118]
exon introduces a stop codon						
GLIAL GROWTH FACTOR		neuron specific	L12259	B	GTGGAGAGATGCCTACTGGCTGCTATTTCAGTCAGCTCTAAGGGAGTGATCAAGGTATGTGGTC CACTTGAATCACG	[102]
exons introducing a phosphorylation site						
H tyrosine hydroxylase, HTH-2		neuron specific immunocytochemistry	M18115	H	GGTAAGAGGGCAGgttaggtgcc	[119], [120], [121], [122]



#### **alternative 3' and polyadenylation**



**pairwise mutually exclusive**

GO alpha, exon 7B		mainly in brain, less in heart and lung, Northern blot	M60161	H	cccccccaacttgttcgcag@AACCGATGCGCAAATCCCTGAAGCTTTGACAGCATCTGCCAAC AACAAATAGTGTCAACAGACAGTCATCATTCTTCTAACAAAGAACATAATTGAGAGAA GATCAAGAAGTCCCCGCTCACCATCTGCTTCTGAATATCACGtagagacccc	[131],
GO alpha, exon 8B		mainly in brain, less in heart and lung, Northern blot	M60162	H	tgccttgcgttttgtctgcag@GCCAGCGCTTACAGAGCGCTGGCTTACATCAGGCCAG TACGAGAGCAAAAGACTGAGCCCCAAAGAGATCTCACGGCACCTGCACCTGCCACGGACAC CAACAACTTCAAGGTTGCTTGTGATGCTGAGCGACGTCTACATCGCCAAAACCTCGGGCT GTGGACACTACTGAGGCCAGCGCCGCTGCCGGCACCTTGCCCTGCCCTGGCTGCCGGCCCC CTCCCCCTGGAAACCAAGCTCCACAGGTCTCA	[131],
GO alpha, exon 7A		mainly in brain, less in heart and lung, Northern blot	M60163	H	tccctttctgcggccgcag@AACCGATGCGCAAATCCCTGAAGCTCCATCTGAAAC AACAAATAGTGTCACTGATCACTCCATCATTCTTCTAACAAAGAACATACTTGGCGAGAA GATCAAGAAGTCTCTTGCACCATCTGCTTCTGAATATCACAGGTTgggtgcga	[131],
GO alpha, exon 8A		mainly in brain, less in heart and lung, Northern blot	M60164	H	tcttcgtgtttgttacag@GCCAAATACCTATGAAAGACGCGAGCCCTACATCAGCACAA GAATAACATCCAGGTTGCTGCTGAGCGCGCTAACAGCACATCATTGCAACACCTCGGGGCT GGGGCTGTACTGACCTTGTGCTGTACCTGAAACCTTATTGtaatgattccagcacccacag aacaaacttcatatccacccatcacacac	[131],

(Table 1. *Cont.*)



### *Distant coordinated expression*

Agrin, exon A		only in neurons exon A and exon B simultaneously used PCR analysis	M97371	C	GTAAATCCCGTAAG	[132]
Agrin, exon B		only in neurons exon A and exon B simultaneously used PCR analysis	M94271	C	TGCCCGACGCATGGACTACCCCTGCTTGAGCCCAG	[132]



FMRF-2, retained intron	regional	neuron immunocytochemistry	specific,	M14958	A	GTTTCCGGAAAAGGTTATGCGATTGGAAAGGGCTCCAGCAATGATGATGAAAGTGGTGATGAC GATGTCGAGGATCTGACGGATAATTGGATGTTCTGGGGGGAAGGGGAAGTAAATAACGGTT CATGAG	[1133 11]
----------------------------	----------	-------------------------------	-----------	--------	---	--	--------------

gene structure unknown

(Table 1. Cont.)

Neurexin I alpha		neuron specific, in situ, protection	M96374	R	CACTCAGGCATGGACACGCTATG	[140]
Neurexin I alpha		neuron specific, in situ, protection	M96374	R	ATTGTATCAGGATTAATGTAATTCCA	[140]
Neurexin I alpha		neuron specific, in situ, protection	M96374	R	GAAACATGATAACGAGCCTGGCGATTCGCTAGACAGCGAATTCCATATCGACTTGGTCGAGTA	[140]
Neurexin I alpha		neuron specific, in situ, protection	M96374	R	GTTGATGAATGGCTACTCGACAAG	[140]
Drebrin	developmental	neuron specific immunocytochemistry	M36961	C	GGCAGCCAGTCCGACTACCGAAAGGTTGGCAGCGGGCTGGAGCCCTGGAGTCAGCCGG CCTCACGCCGCTGGCGAGCAGCAGCGCACCGCCCGCCGAAGAGACGCCAACGCCAA	[141]
Calcineurin A, PP2B α2		brain specific cDNA cloning	M29550	H	GGATGTTCTAGGCCTGAAGACATATCGATTAATCCTCACATAATTTAATGAG	[142]
Calcineurin, PP2B α 1	regional	brain specific, also expressed in thymus	J05480	M	GGCTACTGTTGAGGCTATCGAGGCTGATGAA	[143]

Regional distribution indicates the alternative splicing is different in the various regions of the brain. Developmental regulation indicates a regulation during embryonic and early postnatal development. Experimental evidence lists the experimental methods employed to determine the specificity (in situ: RNA *in situ* hybridisation). If the alternative spliced exons were found to a lesser extent in other tissues, it is indicated under 'experimental evidence'. A '@' sign in the sequences indicates the beginning of a codon. The species are abbreviated as follows: R: rat, M: mouse, H: human, D: drosophila, B: bovine, C: chicken, F: fish. The accession number refers to GenBank. Exon sequences are indicated by capital letters, intron sequences by small letters.

program (NCBI) [24] to find related sequences and entry errors. Vertebrate internal exons and their corresponding downstream donor sites were obtained from the GeneID-datasets [25]. The vertebrate acceptor sites were kindly provided by Dr Knudsen (CEDB, West Florida). In order to produce scores for the combined 5' and 3' splice sites, we randomly selected 200 Genbank (80.0, 12/10/93) vertebrate splice site pairs that flank constitutively spliced exons. cDNA sequences upstream and downstream of the alternatively spliced exons were extracted from Genbank as control sequences for constitutively spliced exons.

LOG-ODD scores were calculated according to Zhang and Marr [26] using the GeneID dataset and randomly extracted splice sites as a comparison. The scoring function is defined as:

$$S_i(X) = \log_2 \frac{P_i(X)}{Q(X)}$$

$P_i(X)$  is the frequency of finding X at position  $i$  that is equal to  $C_i(X)/D_i$ . The normalization  $D_i$  is the sum of the counts  $C_i(X)$  over X (=A,C,G,T). We used  $Q(X)=1/4$  for all X as the random background frequency. The score for a splice site is the sum of the scores for each individual nucleotide.

## RESULTS AND DISCUSSION

### Different classes of alternatively spliced exons in neurons

The different classes of alternatively spliced exons are shown in Table 1. They were arranged according to their gene structure, gene expression and splicing pattern. The splicing pattern is schematically indicated on top of each class. Exons that introduce known regulatory elements like phosphorylation sites, stop codons or frameshifts are listed at the end of each list. The reading frame is indicated by a '@'. The regulation of the alternative splicing, the expression pattern and the experimental techniques used to analyse the alternative exons are included if such information was available in the literature.

### Splice site analysis

The nucleotide composition of splice sites surrounding alternatively spliced exons has been shown to be important for their alternative use [27]. The deviation of splice sites from the consensus sequence [28] seems to decrease binding of splicing factors around the alternatively spliced exon and to limit its use. Mutations in the 5' splice site that interfere with U1 snRNA





Figure 1. Nucleotide usage at the 5' splice site. The percent nucleotide usage of constitutively (left) and neuron specific alternatively spliced exons (right) is pairwise compared. The U1 sequence and the vertebrate consensus splice site sequence are shown at the bottom. The different nucleotides are indicated by different patterns, as indicated in the figure.

binding have been shown to decrease the usage of exons [29, 30]. The composition of the 3' splice site [31] and the branch point has been shown to be important. However, the splice sites of alternative exons do not always deviate from the consensus. For example, the splice sites surrounding the exons coding for substance P [16], sexlethal and the Drosophila P element [8] match the consensus sequence.

### General 5' splice site composition

The 5' splice site consensus sequence reflects base pairing with U1 snRNA. A comparison of 5' splice sites from common and neuron specific alternatively spliced exons is shown in Figure 1. On average, neuron specific 5' splice sites deviate most from the consensus at the +3 and -4 position, where the consensus nucleotide is present 40% less often than in constitutive exons. In position +2, +1, and -3 use of the consensus nucleotide is increased by 8%, at the position -5 it is decreased by 16% and at -6 it is increased by 26%. Addition of these percentages shows that neuron specific 5' splice sites overall use 49% fewer consensus nucleotides complementary to U1 snRNA. However, this deviation is not equally distributed. Most deviation takes place at two positions, +3 (consensus: C) and -4 (consensus: A). Whether this uneven distribution has functional significance has

to be determined. The nucleotide at the -4 position has been postulated to interact with A<sub>49</sub> of the U6 snRNA in a late step in splicing [32, 33]. Compared to constitutive 5' splice sites, alternatively spliced neuronal 5' splice sites have fewer A, but more U at this position, thus on average binding of their 5' splice sites with U6 snRNA might be facilitated. It is interesting to note



**Figure 2.** Nucleotide usage at the 3' splice site. The percent nucleotide usage of constitutively (left) and neuron specific alternatively spliced exons (right) is pairwise compared. The vertebrate consensus sequence is indicated at the bottom; exon sequences are in capital letters. Note the use of A in alternatively spliced exons at the -3 position. The different nucleotides are indicated using the same patterns as in Figure 1.


that the consensus nucleotide at the -4 position in yeast is also U, but the significance of this similarity is unclear.

### General 3' splice site composition

The comparison of the 3' splice sites from neuron specific alternatively spliced exons and common exons reveals differences at the -3 position (Figure 2). Twenty one percent of all neuron-specific exons use an A at this position, compared to only 4% in constitutively used 3' splice sites. The reason why a pyrimidine is conserved at the -3 position is not clear. However, it has been demonstrated that a C→A mutation at this position reduces splicing efficiency by 70% [30]. In addition, in each but the -14 position, neuron specific 3' splice sites have a higher purine content when compared with constitutive 3' splice sites.

### Evaluation of splice sites using LOG-ODD scores


In order to assess the splice site quality, we calculated LOG-ODD scores for each splice site. The LOG-ODD scores express the coincidence of a splice site with the consensus sequence; a higher coincidence generates a higher score. We first calculated a score for each alternative splice site and compared the distribution of individual scores with constitutively spliced exons. Since it has been postulated that the splice sites surrounding an exon define its borders in a concerted way [34], we then calculated the LOG-ODD scores for the splice sites



**Figure 3.** Distribution of splice sites scores in neuron specific and constitutively spliced vertebrate exons. The scores are plotted on the x-axis, the higher the score the better is the match to the consensus sequence. The number of splice sites that accompany a certain score are plotted on the y-axis. Neuron-specific exons are shaded. A. Distribution of neuron-specific 5' splice sites. B. Distribution of common 5' splice sites. A perfect match to U1 would have a score of 12.2. The mean of the distribution is 7.1 for neuron specific exons and 8.0 for constitutively spliced exons. C. Distribution of neuron-specific 3' splice sites. D. Distribution of common 3' splice sites. A 'perfect' 3' splice site (u)<sub>11</sub>cagG would have a score of 15.6. The mean of the distribution is 5.7 for neuron specific exons and 9.4 for constitutively spliced exons. E. Distribution of combined scores of 5' and 3' splice sites that surround a common exon. An exon surrounded by a 'perfect' 3' and 5' splice site would have a score of 27.8. The mean of the distribution is 12.2 for neuron specific exons and 16.9 for constitutively spliced exons.

surrounding one exon. Compared to the control splice sites, the distribution of alternative donor and acceptor splice sites is broader and the mean of the distribution is at a lower score (Figure 3, A and B). This could indicate the presence of two subclasses of splice sites in alternatively spliced exons, with one subclass having the splice sites in consensus and the other subclass having sub-optimal splice sites. When the splice sites surrounding one individual exon are considered, the presence of two subclasses becomes more apparent (Figure 3, E and F). One group of

alternatively spliced exons has splice sites that are sub-optimal, whereas another group has splice sites that score similar to the splice sites of constitutively spliced exons. This could mean that the first group of exons is most likely regulated by their splice site quality, which is in agreement with the exon definition model [34]. Skipping of these exons is the most likely default mechanism. The second group is most likely regulated by other elements like steric hindrance [15], secondary structures [16] or factors that bind to flanking intron sequences [35].



**Figure 4.** Trinucleotide and amino acid distribution in neuron specific (shaded) and constitutively spliced exons (open). **A.** Distribution of overlapping 3-tuples in alternatively and constitutively spliced exons. The actual nucleotide distribution of alternatively and constitutively exons was used to calculate the random nucleotide distribution by random shuffling, the means and the double standard deviations of these random samples are plotted as dots and error bars. Columns indicate the actual percent usage of each 3-tuple. **B.** Distribution of amino acids in alternatively and constitutively spliced exons. The percent usage of each amino acid is indicated. The error bars indicate the standard deviation.

## Nucleotide composition of alternative spliced exons and possible motifs

We analysed the nucleotide composition of the alternatively spliced exons to assess its possible role in exon usage. Overall, alternatively spliced exons are more AU-rich (26% A, 27% C, 24% G, 23% U) than common exons (25% A, 28% C, 27% G, 20% U). The p-values [36] calculated for the differences of the A, C, G, and U values are 0.015, 0.106,  $2.97 \times 10^{-12}$  and  $4.68 \times 10^{-11}$ , respectively. Therefore, the null hypothesis that the nucleotide composition in constitutively and alternatively spliced exons are the same should be rejected at the 0.05 significance level for all nucleotides except C. We conclude from this statistical analysis that the differences in nucleotide composition are significant.

Since several groups reported the involvement of exon motifs in recognition of alternatively spliced exons [13, 37–47] we analysed the neuron specific exon database for k-tuple frequencies according to Claverie *et al.* [48]. The overlapping 3-tuple distributions in both alternatively and constitutively spliced exons are plotted in Figure 4, A. The expected frequencies and the dispersions for random sequences with the same nucleotide compositions were calculated by randomizing each data set and are indicated as error bars in Figure 4, A. The general feature for all exons is the rare use of TA and CG dinucleotides as shown by the deviation below the random expectations, possibly to avoid termination and methylation. Assuming that the nucleotides assemble independently of each other, most of the differences in 3-tuple frequencies between the two exon data sets can be explained by the single nucleotide composition, because the 3-tuple frequencies follow the trend of the random expectations. However, some of the major differences cannot simply be attributed the single nucleotide composition, for example the relative high frequency of trinucleotides AAA and TTT in alternatively spliced exons cannot be explained by the single nucleotide composition and by the amino acid composition of the alternatively spliced exons (Figure 4, B).

In common exons, the trinucleotides CAG, CTG, AGA, TGG and GAG are more frequent than what would be expected in a random assembly. Their high frequency cannot be explained by the amino acid composition of the common exons (Figure 4, B), because only the amino acids F, S, W, Y could contribute to a 3-tuple frequency difference but except F these differences are not correlated to the different trinucleotide distribution. We therefore assume that constraints that lead to the different 3-tuple distribution are due to regulatory requirements on the RNA level. We furthermore conclude that the different 3-tuple composition of alternatively and constitutively spliced exons might have regulatory significance in splicing.

The high frequency of certain trinucleotides probably represents the general characteristics of common exons. In order to see how these 3-tuples are distributed in alternatively spliced exons, we computed the 3-tuple frequency per sequence for the neuron specific exons (each 3-tuple counts at most once in each exon), CAG and AGA were found to be the most frequent, with more than 89% of the alternatively spliced exons containing either CAG or AGA (data not shown).

In order to find motifs common in alternatively spliced exons, we used the RTide program which was designed to search short motifs in multi-sequences [49]. We were unable to identify a motif in the exon sequences or in subset of these sequences. However, our RTide analysis indicated that CAGA might be part of possible motifs in alternatively spliced exons.

The AU rich nucleotide composition and the distribution of tri and tetranucleotides in alternatively spliced exons is in contrast to several AG rich sequence motifs that have been described as necessary for alternative exon usage [38–40, 42, 43, 45–47]. It is tempting to speculate that the AG rich motifs that are in an AU rich context of the alternatively spliced exons serve as signals for the splicing machinery, presumably through binding to an hnRNP. In the light of our analysis, the mutation of these motifs leads to skipping of the alternative exon, because different or no trans factors are now binding to the mutant exon and make it recognizable to the splicing machinery.

Since rapid progress is being made in sequencing and identifying neuron specific exons, we hope that in future updates of this sequence comparison putative motifs will become more clear.

## Update

Since we are planning to update this compilation in the future, we would be thankful for the communication of new or here omitted neuron specific exons. Furthermore, the sequences can be obtained electronically either from stamm@cshl.org or by anonymous ftp from phage.cshl.org in the /pub/science/alt\_exon directory.

## ACKNOWLEDGEMENTS

We thank D.Horowitz, A.R.Krainer, G.J.Mulligan, R.Fislage, D.Casper, C.M.Craft and R.McCombie for helpful discussions and B.Peters for help with computers. This work was supported by an Ausbildungsstipendium of the Deutsche Forschungsgemeinschaft # Sta 399/1-1 to SS, NIH grant RO1-GM43049 to DMH, NIH grant 1KO1 HG00010-02 to MQZ, 1RO1 HG00203-01A1 and DOE # DE-FG02-91ER61190 to TGM. DMH is an Established Investigator of the American Heart Association.

## REFERENCES

- Chikaraishi, D.M., Deeb, S.S., and Sueoka, N. (1978) Cell 13: 111–120.
- Chaudhari, N. and Hahn, W.E. (1983) Science 220: 924–928.
- Milner, R.J. and Sutcliffe, J.G. (1983) Nucl. Acids Res. 11: 5497–5520.
- Ozawa, H., Kushiya, E., and Takahashi, Y. (1980) Bull. Jap. Neurochem. Soc. 19: 338–341.
- Takahashi, Y. (1992) Prog. Neurobiol. 38: 523–569.
- Sutcliffe, J.G. (1988) Ann. Rev. Neurosci. 11: 157–198.
- Sommer, B., Köhler, M., Sprengel, R., and Seuberg, P.H. (1991) Cell 67: 11–19.
- Bingham, P.M., Chou, T.-B., Mims, I., and Zachar, Z. (1988) Trends Genet. 4: 134–138.
- Sommer, B., Keinänen, K., Verdoorn, T.A., Wisden, W., Burnashev, N., Herb, A., Köhler, M., Takagi, T., Sakman, B., and Seuberg, P.H. (1990) Science 249: 1580–1585.
- Hollmann, M., Boulter, J., Maron, C., Beasley, L., Sullivan, J., Pecht, G., and Heinemann, S. (1993) Neuron 10: 943–954.
- Stamm, S., Casper, D., Dinsmore, J., Kaufmann, C.A., Brosius, J., and Helfman, D. (1992) Nucl. Acids Res. 20: 5097–5103.
- Roesser, J.R., Liitschwager, K., and Leff, S.E. (1993) J. Biol. Chem. 268: 8366–8375.
- Stamm, S., Casper, D., and Helfman, D.M. (1993) Soc. Neurosci. Abstr. 19, Part 2, 1699.
- Black, D.L. (1992) Cell 69: 795–807.
- Black, D.L. (1991) Genes & Dev. 5: 389–402.
- Carter, M.S. and Krause, J.E. (1990) J. Neurosci. 10: 2203–2214.
- Zahler, A.M., Neugebauer, K.M., Lane, W.S., and M.B., R. (1993) Science 260: 219–222.
- Branlant, C., Krol, A., Ebel, J.-P., Lazar, E., Haendler, B., and Jacob, M. (1982) EMBO J. 1: 1259–1265.
- McAllister, G., Amara, S.G., and Lerner, M.R. (1988) Proc. Natl. Acad. Sci. USA 85: 5296–5300.

20. Ozcelik, T., Leff, S., Robinson, W., Donlon, T., Lalande, M., Sanjines, E., Schnizel, A., and Francke, U. (1993) *Nat. Genet.* 2: 265–269.
21. Delsert, C.D. and Rosenfeld, M.G. (1992) *J. Biol. Chem.* 267: 14573–14579.
22. Izzo, J.A., Sherman, C.A., and Kusiak, J.W. (1993) *Soc. Neurosci. Abstr.* 19, part 2: 1122.
23. Lees-Miller, J.P., Goodwin, L.O., and Helfman, D.M. (1990) *Mol. Cell Biol.* 10: 1729–1742.
24. Altschul, Stephen, F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990) *J. Mol. Biol.* 215: 403–410.
25. Guigó, R., Knudsen, S., Drake, N., and Smith, T. (1992) *J. Mol. Biol.* 226: 141–157.
26. Zhang, M.Q. and Marr, T.G. (1994) *Nucl. Acids Res.* 22: 1750–1759.
27. Horowitz, D.S. and Krainer, A.R. (1994) *Trends Genet.* 10: 100–106.
28. Stephens, R.M. and Schneider, T.D. (1992) *J. Mol. Biol.* 228: 1124–1136.
29. Kuo, H.-C., Nasim, F.-U.H., and Grabowski, P.J. (1991) *Science* 251: 1045–1050.
30. Aeby, M., Hornig, H., Padgett, R.A., Reiser, J., and Weissmann, C. (1986) *Cell* 47: 555–565.
31. Adema, G.J., Hulst, K.L., and Baas, P.D. (1990) *Nucl. Acids Res.* 18: 5365–5373.
32. Sontheimer, E.J. and Steitz, J.A. (1993) *Science* 262: 1989–1996.
33. Lesser, C.F. and Guthrie, C. (1993) *Science* 262: 1982–1988.
34. Robberson, B.L., Cote, G.J., and Berget, S.M. (1990) *Mol. Cell. Biol.* 10: 84–94.
35. Mulligan, G.J., Guo, W., Wormsley, S., and Helfman, D.M. (1992) *J. Biol. Chem.* 267: 25480–25487.
36. Fleiss, J.L. *Statistical methods for rates and proportions*. 2nd ed ed. 1981 New York: Wiley.
37. Katz, R.A. and Skalka, A.M. (1990) *Mol. Cell. Biol.* 10: 696–704.
38. Mardon, H.J., Sebastio, G., and Baralle, F.E. (1987) *Nucl. Acids Res.* 15: 7725–7733.
39. Watakabe, A., Tanaka, K., and Shimura, Y. (1993) *Genes and Dev.* 7: 407–418.
40. Hampson, R.K., LaFollette, L., and Rottman, F.M. (1989) *Mol. Cell. Biol.* 9: 1604–1610.
41. Cooper, T.A. (1992) *J. Biol. Chem.* 267: 5330–5338.
42. Steingrimsdottir, H.G., Rowley, G., Dorado, G., Cole, J., and Lehmann, A.R. (1992) *Nucl. Acids Res.* 20: 1201–1208.
43. Fu, X.-D., Kats, R.A., Skalka, A.M., and Maniatis, T. (1991) *Genes and Dev.* 5: 211–220.
44. Yeakley, J.M., Hedjran, F., Morfin, J.P., Merillat, N., Rosenfeld, M.G., and Emeson, R.B. (1993) *Mol. Cell. Biol.* 13: 5999–6011.
45. Guo, W., Mulligan, G.J., Wormsley, S., and Helfman, D.M. (1991) *Genes Dev.* 5: 2096–2107.
46. Lavigne, A., La Branche, H., Kornblith, A.R., and Chabot, B. (1993) *Genes and Dev.* 7: 2405–2417.
47. Xu, R., Teng, J., and Cooper, T.A. (1993) *Mol. Cell. Biol.* 13: 3660–3674.
48. Claverie, J.-M., Sauvaget, I., and Bougueret, L. (1990) *Meth. Enzymol.* 183: 237–252.
49. Galas, D.J., Waterman, M.S., and Eggert, M. (1985) *J. Mol. Biol.* 186: 117–128.
50. Kuhse, J., Kuryatov, A., Maulet, Y., Malosio, M.L., Schmieden, V., and Betz, H. (1991) *FEBS letters* 283: 73–77.
51. Monyer, H., Seuberg, P.H., and Wisden, W. (1991) *Neuron* 6: 799–810.
52. Snutch, T.P., Tomlinson, W.J., Leonard, J.P., and Gilbert, M.M. (1991) *Neuron* 7: 45–57.
53. Ma, W.-J., Holz, R.W., and Uhler, M.D. (1992) *J. Biol. Chem.* 267: 22728–22732.
54. Buck, L.B., Bigelow, J.M., and Axel, R. (1987) *Cell* 51: 127–133.
55. Kirchhausen, T., Scarmato, P., Harrison, S.C., Monroe, J.J., Chow, P.E., Mattaliano, R.J., Ramachandran, K.L., Smart, J.E., Ahn, A.H., and Brosius, J. (1987) *Science* 236: 320–324.
56. Jackson, A.P. and Parham, P. (1988) *J. Biol. Chem.* 263: 16688–16695.
57. Stamm, S. . unpublished
58. Maycox, P.R., Link, E., Reetz, A., Morris, S.A., and Jahn, R. (1992) *J. Cell Biol.* 118: 1379–1388.
59. Jackson, A.P., Seow, H.F., Holmes, N., Drickamer, K., and Parham, P. (1987) *Nature* 326: 154–159.
60. Owens, G.C., Edelman, G.M., and Cunningham, B.A. (1987) *Proc. Natl. Acad. Sci.* 84: 294–298.
61. Small, S.J. and Akeson, R. (1990) *J. Cell Biol.* 111: 2089–2096.
62. Small, S.J., Haines, S.L., and Akeson, R.A. (1988) *Neuron* 1: 1007–1017.
63. Santoni, M.J., Barthels, D., Vopper, G., Boned, A., Goridis, C., and Wille, W. (1989) *EMBO J.* 8: 385–392.
64. Huang, J.P., Tang, C.J., Kou, G.-H., Marchesi, V.T., and Benz, E.J. (1993) *J. Biol. Chem.* 268: 3758–3766.
65. Levy, J.B., Dorai, T., Wang, L.-H., and Brugge, J.S. (1987) *Mol. Cell. Biol.* 7: 4142–4145.
66. Raulf, F., Robertson, S.M., and Schartl, M. (1989) *J. Neurosci. Res.* 24: 81–88.
67. Pyper, J.M. and Bolen, J.B. (1989) *J. Neurosci. Res.* 24: 89–96.
68. Collett, J.W. and Steele, R.E. (1992) *Dev. Biol.* 152: 194–198.
69. Morgan, B.A., Johnson, W.A., and Hirsh, J. (1986) *EMBO J.* 5: 3335–3342.
70. Nawa, H., Kotani, H., and Nakanishi, S. (1984) *Nature* 312: 729–734.
71. Prior, P., Schmitt, B., Grenningloh, G., Pribilla, I., Multhaup, G., Langosch, D., Kirsch, J., and Betz, H. (1992) *Neuron* 8: 1161–1170.
72. Cyr, J.L., Pfister, K.K., Bloom, G.S., Slaughter, C.A., and Brady, S.T. (1991) *Proc. Natl. Acad. Sci. USA* 88: 10114–10118.
73. Danoff, S.K., Ferris, C.D., Donath, C., Fischer, G.A., Munemitsu, S., Ullrich, A., Snyder, S.H., and Ross, C.A. (1991) *Proc. Natl. Acad. Sci. USA* 88: 2951–2955.
74. Fujita, N., Sato, S., Kurihara, T., Inuzuka, T., Takahashi, Y., and Miyatake, T. (1988) *FEBS Lett.* 232: 323–327.
75. Lai, C., Brow, M.A., Nave, K.A., Noronha, A.B., Quarles, R.H., Bloom, F.E., Milner, R.J., and Sutcliffe, J.G. (1987) *Proc. Natl. Acad. Sci. USA* 84: 4337–4341.
76. Staugaitis, S.M., Smith, P.R., and Colman, D.R. (1990) *J. Cell. Biol.* 110: 1719–1727.
77. Fujita, N., Sato, S., Kurihara, T., Kuwano, R., Sakimura, K., Inuzuka, T., Takahashi, Y., and Miyatake, T. (1989) *Biochem. Biophys. Res. Commun.* 165: 1162–1169.
78. Pyper, J. and Bolen, J.B. (1990) *Mol. Cell Biol.* 10: 2035–2040.
79. Ono, Y., Kikkawa, U., Ogita, K., Fujii, T., Kurokawa, T., Asaoka, Y., Sekiguchi, K., Ase, K., Igarashi, K., and Nishizuka, Y. (1987) *Science* 236: 1116–1120.
80. Hirata, M., Saito, N., Kono, M., and Tanaka, C. (1991) *Dev. Brain Res.* 62: 229–238.
81. Coussens, L., Rhee, L., Parker, P.J., and Ullrich, A. (1987) *DNA* 6: 389–394.
82. Wirth, T., Priess, A., Annweiler, A., Zwilling, S., and Oeler, B. (1991) *Nucl. Acids Res.* 19: 43–51.
83. Lillycrop, K.A. and Latchman, D.S. (1992) *J. Biol. Chem.* 267: 24960–24965.
84. Giros, B., Martres, M.-P., Pilon, C., Sokoloff, P., and Schwartz, J.-C. (1991) *Biochem. Biophys. Res. Com.* 176: 1584–1592.
85. Nakanishi, N., Axel, R., and Schneider, N.A. (1992) *Proc. Natl. Acad. Sci. USA* 89: 8552–8556.
86. Giros, B., Sokoloff, P., Riou, J.-F., Emorine, L.J., and Schwartz, J.-C. (1989) *Nature* 342: 923–926.
87. Monsma, J.F.J., McVittie, L.D., Gersen, C.R., Mahan, L.C., and Sibley, D.R. (1989) *Nature* 342: 926–929.
88. Miller, J.C., Wang, Y., and Filer, D. (1990) *Biochem. Biophys. Res. Comm.* 166: 109–112.
89. Dal Toso, R., Sommer, B., Ewert, M., Herb, A., Prichett, D.B., Bach, A., Shivers, B.D., and Seuberg, P.H. (1989) *EMBO J.* 8: 4025–4034.
90. Selbie, L.A., Hayes, G., and Shine, J. (1989) *DNA* 8: 683–689.
91. Montmayeur, J.P., Bausero, P., Amlaiky, N., Maroteaux, L., Hen, R., and Borrelli, E. (1991) *FEBS lett.* 278: 239–243.
92. Schaller, K.L., Krzemien, D.M., McKenna, N.M., and Caldwell, J.H. (1992) *J. Neurosci.* 12: 1370–1381.
93. Andreatis, A., Brown, W.M., and Kosik, K.S. (1992) *Biochemistry* 31: 10626–10633.
94. Goedert, M., Spillantini, M.G., Jakes, R., Rutherford, D., and Crowther, R.A. (1989) *Neuron* 3: 519–526.
95. Goedert, M., Spillantini, M.G., Potier, M.C., Ulrich, J., and Crowther, R.A. (1989) *EMBO J.* 8: 393–399.
96. Goedert, M., Spillantini, M.G., and Crowther, R.A. (1992) *Proc. Natl. Acad. Sci. USA* 89: 1983–1987.
97. Himmeler, A. (1989) *Mol. Cell Biol.* 9: 1389–1396.
98. Papandrikopoulou, A., Doll, T., Tucker, R.P., Garner, C.C., and Matus, A. (1989) *Nature* 340: 650–652.
99. Ushkaryov, Y.A. and Südhof, T.C. (1993) *Proc. Natl. Acad. Sci. USA* 90: 6410–6414.
100. Bennett, M.K. and Kennedy, M.B. (1987) *Proc. Natl. Acad. Sci. USA* 84: 1794–1798.
101. Bulleit, R.F., Bennett, M.K., Molloy, S.S., Hurley, J.B., and Kennedy, M.B. (1988) *Neuron* 1: 63–72.
102. Marchionni, M.A., Goodearl, A.D., Chen, M.S., Birmingham, M.O.,

- Kirk, C., Hendricks, M., Danehy, F., Misumi, D., Sudhalter, J., Kobayashi, K., Wroblewski, D., Lynch, C., Baldassare, M., Hiles, I., Davis, J.B., Hsuan, J.J., Totty, N.F., Waterfield, M.D., Stroobant, P., and Gwynne, D. (1993) *Nature* 362: 312–318.
103. Duilio, A., Zambrano, N., Mogavero, A.R., Ammendola, R., Cimino, F., and Russo, T. (1991) *Nucl. Acids Res.* 19: 5269–5274.
104. Lorenzi, M.V., Trinidad, A.C., Zhang, R., and Strauss, W.L. (1992) *DNA* 11: 593–603.
105. Whiting, P., McKernan, R.M., and Iversen, L.L. (1990) *Proc. Natl. Acad. Sci. USA* 87: 9966–9970.
106. Glencorse, T.A., Bateson, A.N., and Darlison, M.G. (1990) *Nucl. Acids Res.* 18: 7157.
107. Kofuji, P., Wang, J.B., Moss, S.J., Huganir, R.L., and Burt, D.R. (1991) *J. Neurochem.* 56: 713–715.
108. Wang, J.-B. and Burt, D.R. (1991) *Brain Res. Bull.* 27: 731–735.
109. Miura, M., Kobayashi, M., Asou, H., and Uyemura, K. (1991) *Febs Lett* 289: 91–95.
110. Malosio, M.L., Grenningloh, G., Kuhse, J., Schmieden, V., Schmitt, B., Prior, P., and H., B. (1991) *J. Biol. Chem.* 266: 2048–2053.
111. Fishburn, C.S., Belleli, D., David, C., and Fuchs, S. (1993) *J. Biol. Chem.* 268: 5872–5878.
112. Oates, E. and Herbert, E. (1984) *J. Biol. Chem.* 259: 7421–7425.
113. Maurer, R.A., Erwin, C.R., and Donelson, J.E. (1981) *J. Biol. Chem.* 256: 10524–10528.
114. Richter, K., Good, P.J., and Dawid, I.G. (1990) *New Biologist* 2: 556–565.
115. Südhof, T.C., Czernik, A.J., Kao, H.-T., Takei, K., Johnston, P.A., Horiuchi, A., Kanazir, S.D., Wagner, M.A., Perin, M.S., De Camilli, P., and Greengard, P. (1989) *Science* 245: 1474–1480.
116. Südhof, T.C. (1990) *J. Biol. Chem.* 265: 7849–7852.
117. Bateson, A.N., Lasham, A., and Darlison, M.G. (1991) *J. Neurochem.* 56: 1437–1440.
118. Yokoyama, S., Imoto, K., Kawamura, T., Higashida, H., Iwabe, N., Miyata, T., and Numa, S. (1989) *FEBS Lett.* 259: 37–42.
119. Grima, B., Lamouroux, A., Boni, C., Julien, J.F., Javoy, A.F., and Mallet, J. (1987) *Nature* 326: 707–711.
120. O’Malley, K.L., Anhalt, M.J., Martin, B.M., Kelsoe, J.R., Winfield, S.L., and Ginns, E.I. (1987) *Biochemistry* 26: 6910–6914.
121. Le Bourdellès, B., Horellou, P., Le Caer, J.-P., Denèfle, P., Latta, M., Haavik, J., Guibert, B., Mayaux, J.-F., and Mallet, J. (1991) *J. Biol. Chem.* 266: 17124–17130.
122. Kaneda, N., Kobayashi, K., Ichinose, H., Kishi, F., Nakazawa, A., Kurosawa, Y., Fujita, K., and Nagatsu, T. (1987) *Biochem. Biophys. Res. Commun.* 146: 971–975.
123. Stamm, S., Casper, D., Lees-Miller, J.P., and Helfman, D.M. (1993) *Proc. Natl. Acad. Sci. USA* 90: 9857–9861.
124. Forry-Schaudies, S. and Hughes, S.H. (1991) *J. Biol. Chem.* 266: 13821–13827.
125. Hortsch, M., Bieber, A.J., Patel, N.H., and Goodman, C.S. (1990) *Neuron* 4: 467–709.
126. Amara, S., Evans, R.M., and Rosenfeld, M.G. (1984) *Mol. Cell Biol.* 4: 2151–2160.
127. Amara, S.G., Jonas, V., Rosenfeld, M.G., Ong, E.S., and Evans, R.M. (1982) *Nature* 298: 240–244.
128. Leff, S.E., Evans, R.M., and Rosenfeld, M.G. (1987) *Cell* 48: 517–524.
129. Chang, J.-H. and Olson, M.O.J. (1989) *J. Biol. Chem.* 264: 11732–11737.
130. Luneau, C.J., Williams, J.B., Marshall, J., Levitan, E.S., Oliva, C., Smith, J., Antanavage, J., Folander, K., Stein, R.B., Swanson, R., Kaczmarek, L.K., and Buhrow, S.A. (1991) *Proc. Natl. Acad. Sci.* 88: 3932–3936.
131. Tsukamoto, T., Toyama, R., Itoh, H., Kozasa, T., Matsuoaka, M., and Kaziro, Y. (1991) *Proc. Natl. Acad. Sci. USA* 88: 2974–2978.
132. Ruegg, M., Tsim, K.W., Horton, S.E., Kröger, S., Escher, G., Gensch, E.M., and McMahan, U.J. (1992) *Neuron* 8: 691–699.
133. Schaefer, M., Picciotto, M.R., Kreiner, T., Kaldany, R.-R., Taussig, R., and Scheller, R. (1985) *Cell* 41: 457–467.
134. Hui, A., Ellinor, P.T., Krizanova, O., Wang, J.-J., Diebold, R.J., and Schwartz, A. (1991) *Neuron* 7: 35–44.
135. Kim, H.-L., Kim, H., Lee, P., King, R.G., and Chin, H. (1992) *Proc. Natl. Acad. Sci. USA* 89: 3251–3255.
136. Rettig, J., Wunder, F., Stocker, M., Lichtenhagen, R., Mastiaux, F., Beckh, S., Kues, W., Pedarzani, P., Schröter, K.H., Ruppertsberg, J.P., Veh, R., and Pongs, O. (1992) *EMBO J.* 11: 2473–2486.
137. McCormack, T., Vega-Saenz de Mira, E.C., and Rudy, B. (1991) *Proc. Natl. Acad. Sci. USA* 88: 4060.
138. Luneau, C., Wiedmann, R., Smith, J.S., and Williams, J.B. (1991) *FEBS lett.* 288: 163–167.
139. Goldman, D., Deneris, E., Luyten, W., Kochhar, A., Patrick, J., and Heinemann, S. (1987) *Cell* 48: 965–973.
140. Ushkaryov, Y.A., Petrenko, A.G., Geppert, M., and Südhof, T.C. (1992) *Science* 257: 50–56.
141. Kojima, N., Kato, Y., Shirao, T., and Obata, K. (1988) *Mol. Brain Res.* 4: 207–215.
142. Guerini, D. and Klee, C.B. (1989) *Proc. Natl. Acad. Sci. USA* 86: 9183–9187.
143. Kincaid, R.L., Giri, P.R., Higuchi, S., Tamura, J., Dixon, S.C., Marietta, C.A., Amorese, D.A., and Martin, B.M. (1991) *J. Biol. Chem.* 265: 11312–11319.