

Sex-Specific Stress-Related Behavioral Phenotypes and Central Amygdala Dysfunction in a Mouse Model of 16p11.2 Microdeletion

Jacqueline Giovanniello, Sandra Ahrens, Kai Yu, and Bo Li

ABSTRACT

BACKGROUND: Substantial evidence indicates that a microdeletion on human chromosome 16p11.2 is linked to neurodevelopmental disorders, including autism spectrum disorder (ASD). Carriers of this deletion show divergent symptoms besides the core features of autism spectrum disorder, such as anxiety and emotional symptoms. The neural mechanisms underlying these symptoms are poorly understood.

METHODS: We used mice heterozygous for a deletion allele of the genomic region corresponding to the human 16p11.2 microdeletion locus (i.e., *16p11.2 del/+* mice) and their sex-matched wild-type littermates for the study and examined their anxiety-related behaviors, auditory perception, and central amygdala circuit function using behavioral, circuit tracing, and electrophysiological techniques.

RESULTS: Mice heterozygous for a deletion allele of the genomic region corresponding to the human 16p11.2 microdeletion locus (i.e., *16p11.2 del/+* mice) had sex-specific anxiety-related behavioral and neural circuit changes. Specifically, we found that female, but not male, *16p11.2 del/+* mice showed enhanced fear generalization—a hallmark of anxiety disorders—after auditory fear conditioning and displayed increased anxiety-like behaviors after physical restraint stress. Notably, such sex-specific behavioral changes were paralleled by an increase in activity in central amygdala neurons projecting to the globus pallidus in female, but not male, *16p11.2 del/+* mice.

CONCLUSIONS: Together, these results reveal female-specific anxiety phenotypes related to 16p11.2 microdeletion syndrome and a potential underlying neural circuit mechanism. Our study therefore identifies previously underappreciated sex-specific behavioral and neural changes in a genetic model of 16p11.2 microdeletion syndrome and highlights the importance of investigating female-specific aspects of this syndrome for targeted treatment strategies.

<https://doi.org/10.1016/j.bpsgos.2021.01.001>

Autism spectrum disorder (ASD) is a spectrum of neurodevelopmental conditions defined by two major diagnostic criteria: “persistent deficits in social communication and social interaction across multiple contexts” and “restricted, repetitive patterns of behavior, interests, or activities” (1). Patients with ASD commonly have one or more comorbid conditions, including intellectual disability (2–4), attention-deficit/hyperactivity disorder (5–8), obsessive-compulsive disorder (9,10), anxiety (11–13), and depression (13–16), and are at increased risk for suicidality, particularly female patients (17–19).

It is well documented that ASD is about 4 times more common in males than in females with an exception for X-linked syndromes, such as Rett syndrome, which is more common in females (20). There is significant evidence of divergence among core symptoms of ASD based on sex. Specifically, many studies have found reduced severity of repetitive and/or stereotyped behaviors in females with ASD compared with males (21–25). In contrast, females show different social impairments than males (22,26–29). These tend

toward more internalizing symptoms and emotional disturbance (30–34). Females with ASD also show increased risk of eating disorders (35), sensory impairments (36), sleep disturbances (37), epilepsy, and learning disorders (38). It has been suggested that females may camouflage their autism phenotypes better than males owing to fewer social impairments and better executive functioning (39) as well as reduced externalizing symptoms (29). One way that emotional phenotypes often manifest is as anxiety disorders. In the general population, females have an increased prevalence of stress-related disorders, such as anxiety, depression, and posttraumatic stress disorder (40–43). Therefore, it is possible that anxiety-like phenotypes may present differently in males and females with ASD.

A major limitation of much of the research in ASD has been its emphasis on males. This is not exclusive to ASD research, as most research is done in male subjects (44). Among neuroscience studies in general, the sex bias of human subjects is approximately 5.5 males for every female and with a ratio much higher among animal studies (45). This bias

precludes our understanding of autism in females and limits our development of effective treatment strategies. Therefore, we sought to examine whether sex differences exist in stress-related behaviors in a mouse model of ASD. To this end, we used a model that mimics a microdeletion on human chromosome 16p11.2. Notably, this deletion is one of the most common genetic variations found in ASD, accounting for approximately 1% of ASD cases (46–51). Patients with this deletion show repetitive behaviors, hyperactivity, intellectual disability, motor and speech/language delay, and anxiety symptoms (52–56). Of note, individuals carrying the 16p11.2 deletion, including carriers without ASD, often receive diagnoses of anxiety disorders or other mood disorders (57).

The mouse model we used was generated by Horev *et al.* (58), and is one of three independently generated mouse genetic models that mimic the 16p11.2 microdeletion (58–60). These models, which were created by deleting largely similar genomic intervals in mouse chromosome 7 corresponding to the syntenic 16p11.2 microdeletion region in humans, exhibit overlapping phenotypes (58–60). In particular, heterozygous deletion mice—hereafter referred to as *16p11.2 del/+* mice—in each of these lines share basic phenotypes, such as low body weight and perinatal mortality, and, importantly, show behavioral phenotypes related to the symptoms of human 16p11.2 microdeletion carriers. These phenotypes include increased locomotor activity, stereotyped and repetitive behaviors, sleep deficits, recognition memory deficits, reward learning deficits, and social deficits (58–66).

A few studies examined *16p11.2 del/+* mice for anxiety- or fear-related behaviors, but with mixed results. When tested in the open field test and elevated plus maze (EPM) test, assays conventionally used to assess anxiety in rodents, these mice do not appear different from wild-type mice (60,61,67) [however, see (68)]. The *16p11.2 del/+* mice were also examined in fear conditioning paradigms. One study shows that *16p11.2 del/+* mice have impaired contextual fear conditioning (69), whereas other studies show that *16p11.2 del/+* mice have normal contextual fear conditioning and normal visually cued fear conditioning (61,67).

Recent studies indicate that environmental factors can exacerbate ASD symptoms and impairments in cognitive and adaptive behaviors in 16p11.2 deletion carriers (70), and *16p11.2 del/+* mice show altered coping in response to stress compared with wild-type littermates (61,71). In light of these findings and studies showing that males and females can exhibit very different behavioral responses to threats or stress (72,73), we reasoned that under a stressful situation *16p11.2 del/+* mice may exhibit sex-specific behavioral changes. However, a potential sex-specific effect of the *16p11.2* deletion on anxiety- or fear-related behaviors in mice has not been considered until recently (67). Furthermore, only simple assays, such as the open field test and EPM test, have been used to assess baseline anxiety in *16p11.2 del/+* mice, which may not be sufficient to reveal potential changes in anxiety or fear processing in response to stress in these mice.

To address these issues, in the current study we examined anxiety-related behaviors under different stress conditions in both male and female *16p11.2 del/+* mice and their wild-type littermates. We found that female, but not male, *16p11.2 del/+* mice showed enhanced fear generalization, a hallmark of anxiety disorders (74), after auditory fear conditioning. Furthermore, although

at baseline *16p11.2 del/+* mice were not different from their wild-type littermates in the EPM test, consistent with previous studies (60,61,67), we found that female, but not male, *16p11.2 del/+* mice showed enhanced anxiety in the EPM test after acute restraint stress. Lastly, we found that these sex-specific behavioral changes were paralleled by an increase in activity in the central amygdala (CeA)—a major limbic structure that regulates anxiety in rodents and primates (75–77)—of female, but not male, *16p11.2 del/+* mice. Together, our work suggests that 16p11.2 microdeletion differentially affects males and females and may disproportionately disrupt brain functions related to stress regulation in females. These findings provide insight into understanding how ASD may present differently in females at behavioral and neuronal levels and raise the question of whether changes to treatment and diagnostic strategies based on sex should be considered.

METHODS AND MATERIALS

Animals

To breed *16p11.2 del/+* mice, we used *16p11.2 del/+* male mice and C57/B6 female mice purchased from the Jackson Laboratory (Bar Harbor, ME) or similar breeders obtained from Pavel Osten's laboratory at Cold Spring Harbor Laboratory. Breeders were housed with a cardboard Bio-Hut (Bio-Serv, Flemington, NJ) under a 12-hour light/dark cycle (7 AM to 7 PM light) with food and water available ad libitum. As *16p11.2 del/+* mice exhibit postnatal lethality (58), in breeding cages only, standard rodent chow (Purina LabDiet, Gray Summit, MO) was supplemented with DietGel Boost (ClearH₂O, Westbrook, ME), a high-calorie liquid diet that increased survival of *16p11.2 del/+* pups. Pups were weaned at 3 weeks of age and group housed with wild-type littermates. Mice were genotyped for 16p11.2 microdeletion between 4 and 8 weeks of age with primers provided by Alea Mills' laboratory at Cold Spring Harbor Laboratory.

Mice 2–4 months old were used for all behavioral experiments. Mice 6–10 weeks old were used for all electrophysiology experiments. All experimental mice were housed under a 12-hour light/dark cycle (7 AM to 7 PM light) in groups of 2–5 animals, containing both *16p11.2 del/+* mice and their wild-type littermates. Food and water were available ad libitum. All behavioral experiments were performed during the light cycle. Littermates were randomly assigned to different groups before experiments. All experimental procedures were approved by the Institutional Animal Care and Use Committee of Cold Spring Harbor Laboratory and performed in accordance with the U.S. National Institutes of Health guidelines.

See Supplemental Methods for details on Behavioral Tasks, Stereotaxic Surgery, and In Vitro Electrophysiology.

Data Analysis and Statistics

Statistical analyses were performed with Prism 7 (GraphPad Software, La Jolla, CA). Normality was tested by D'Agostino-Pearson or Shapiro-Wilk normality tests. Non-normal datasets were log-transformed for normality before statistical testing. All behavioral experiments were controlled by computer systems, and data were collected and analyzed in an automated and unbiased way. Virus-injected animals in which the injection site

was incorrect were excluded. No other animals were excluded. Sex was found to be a significant factor in percent freezing during fear conditioning retrieval ($F_{1,83} = 8.740, p = .0041$, two-way repeated measures analysis of variance) (Figure 1). Given this, in addition to our a priori hypothesis that sex differences would be present based on sex bias in the prevalence of human autism and anxiety disorders, all results have been analyzed separately by sex.

RESULTS

Female-Specific Increase in Fear Generalization in 16p11.2 *del/+* Mice

One hallmark of anxiety disorders is fear generalization (74). Fear generalization can be assessed in mice using a fear conditioning paradigm with a discrimination component (see Supplemental Methods), in which mice are trained to associate one auditory stimulus (conditioned stimulus [CS]) (CS+) with a foot shock (unconditioned stimulus [US]), while a different auditory stimulus (CS-) is presented without the shock. In a fear retrieval test 24 hours following the conditioning, freezing in response to the CS+ and freezing in response to the CS- are measured and used to calculate a discrimination index, which is the difference between an animal's average freezing to the CS+ and to the CS-, normalized to the sum of the two measurements.

Interestingly, we found that during a habituation session before the conditioning, female 16p11.2 *del/+* mice showed a small (10%–20%) but robust increase in freezing to the auditory stimuli compared with their wild-type littermates (Figure 1A, left). Male 16p11.2 *del/+* mice did not show such a change (Figure 1A, right). However, we did not observe a significant difference in freezing during the first tone presentation in the subsequent conditioning session (i.e., before mice received any shocks) between genotypes for either the female or the male mice (Figure 1B, D), suggesting that the enhanced freezing in 16p11.2 *del/+* female mice during habituation may be related to the fact that the auditory stimuli were novel to the animals.

After fear conditioning and on memory retrieval, both female and male 16p11.2 *del/+* mice showed levels of freezing similar to those of their wild-type littermates in response to the CS+ (Figure 1B, D), consistent with previous findings that 16p11.2 *del/+* mice have intact fear conditioning (61,67). Surprisingly, however, female, but not male, 16p11.2 *del/+* mice showed increased freezing to the CS- compared with wild-type littermates (Figure 1B, D), resulting in reduced levels of fear discrimination in female, but not male, 16p11.2 *del/+* animals (Figure 1C, E). In addition, we found that female, but not male, 16p11.2 *del/+* mice showed enhanced reactions to foot shocks compared with wild-type mice, as measured by enhanced movement velocity and distance immediately following shocks of varying intensities (Figure 2). These results suggest that female 16p11.2 *del/+* mice have enhanced fear generalization following fear conditioning, which could result from heightened alertness (as indicated by increased freezing during habituation) and/or an increase in sensitivity to aversive stimuli (as indicated by increase reactivity to foot shocks).

16p11.2 *del/+* Mice Have Normal Auditory Perception

An alternative explanation for the enhanced fear generalization in female 16p11.2 *del/+* mice is that these mice have an impairment in auditory processing, such that they cannot effectively discriminate between a 4-kHz tone and a 12-kHz tone, which were used as CS+ and CS-, respectively, during fear conditioning. To test this possibility, we trained a new cohort of mice, including 16p11.2 *del/+* mice and their wild-type littermates, in an auditory two-alternative choice task that depended on discriminating between a 4-kHz tone and a 12-kHz tone (Figure 3A) (see Supplemental Methods) (78). Both female and male 16p11.2 *del/+* mice learned the two-alternative choice task at a rate similar to that of their wild-type littermates (Figure 3B, C). In fact, male 16p11.2 *del/+* mice tended to be faster than wild-type mice in learning the task (Figure 3C), though this difference did not reach significance. In addition, the performance of female and male 16p11.2 *del/+* mice in discriminating a series of sounds with frequencies ranging from 4 to 12 kHz (Figure 3D–F and H–J), or with different intensities (Figure 3G, K), was indistinguishable from their wild-type littermates. These results indicate that 16p11.2 microdeletion does not affect auditory perception in mice, ruling out the possibility that the enhanced fear generalization in female 16p11.2 *del/+* mice is confounded by an impairment in auditory processing in these mice.

Stress Induces an Increase in Anxiety in Female 16p11.2 *del/+* Mice

In fear conditioning, mice receive electric shocks as the aversive US, which is a significant stressor to animals. Therefore, the enhanced fear generalization in female 16p11.2 *del/+* mice after fear conditioning points to a possibility that these animals are prone to stress-induced anxiety. To further test this possibility, we sought to examine anxiety-like behaviors in mice subjected to a different stressor. For this purpose, we used physical restraint (see Supplemental Methods), a well-characterized stress induction procedure in rodents that has been shown to affect the function of the CeA (79,80). As described previously (81), animals were physically restrained in a well-ventilated 50-mL conical tube for 2 hours in a dark, sound-attenuated box. The animals were tested 24 hours later in the EPM (see Supplemental Methods). We found a significant interaction between sex and genotype in the time spent in the open arms (Figure 4A) and significant effects of sex on movement velocity (Figure 4B) and distance traveled (Figure 4C). Post hoc analysis revealed that the stressed female 16p11.2 *del/+* mice spent significantly less time in the open arms of the EPM compared with their female wild-type littermates (Figure 4A). We did not find any change in time spent in the open arms in male 16p11.2 *del/+* mice.

We also examined anxiety levels in naïve mice using the EPM test. Compared with naïve female or male wild-type littermates, naïve female or male 16p11.2 *del/+* mice did not show any change in the time spent in the open arms (Figure 4D), movement velocity (Figure 4E), or distance traveled (Figure 4F). This




Figure 1. Female *16p11.2 del/+* mice exhibit fear generalization following fear conditioning. **(A)** Freezing behavior of male and female *16p11.2 del/+* mice and their respective WT littermates in response to CS+ and CS- during habituation (female [*16p11.2 del/+*, $n = 15$, WT, $n = 16$]: $F_{1,29} = 6.023$, $p = .0204$; male [*16p11.2 del/+*, $n = 28$, WT, $n = 28$]: $F_{1,54} = 0.3433$, $p = .5604$; * $p < .05$; two-way analysis of variance with repeated measures). **(B)** Freezing to each stimulus presentation during conditioning and retrieval for female mice (conditioning: $F_{1,29} = 1.419$, $p = .2432$; CS+ retrieval: $F_{1,29} = 0.4314$, $p = .5165$; CS- retrieval: $F_{1,29} = 5.765$, $p = .0230$; * $p < .05$; two-way analysis of variance with repeated measures and post hoc Sidak test). **(C)** Discrimination index [(CS+ - CS-)/(CS+ + CS-)] for female mice (* $p = .0192$, Mann-Whitney t test). **(D)** Freezing to each stimulus presentation during conditioning and retrieval for male mice (conditioning: $F_{1,54} = 0.9938$, $p = .3233$; CS+ retrieval: $F_{1,54} = 0.6327$, $p = .4298$; CS- retrieval: $F_{1,54} = 0.8779$, $p = .3530$; two-way analysis of variance with repeated measures). **(E)** Discrimination index for male mice ($p = .3742$, Mann-Whitney t test). Data are presented as mean \pm SEM. CS, conditioned stimulus; n.s., not significant; WT, wild-type.

result is consistent with previous findings (60,61,67). Together, our results indicate that female *16p11.2 del/+* mice have increased susceptibility to stress-induced anxiety.

16p11.2 del/+ Mice Have CeA Dysfunction

Previous studies have revealed that the CeA is particularly responsive to stress and is a major contributor to anxiety-related behaviors (75–77). Therefore, we examined whether the *16p11.2* microdeletion affects CeA neuronal function in a sex-specific manner. We recorded miniature excitatory postsynaptic currents (mEPSCs)—a measurement of total excitatory synaptic drive onto the recorded neurons—from CeA neurons in acute brain slices prepared from female or male *16p11.2 del/+* mice as well as their female or male wild-type littermates (Figure 5A). We found significant effects of sex and genotype on mEPSC frequency in randomly recorded CeA neurons (Figure 5B–D). Post hoc analysis revealed that female mice with *16p11.2* microdeletion specifically had increased mEPSC frequency compared with female wild-type littermates. There was no difference in mEPSC amplitude between genotypes or sexes (Figure 5E). These results indicate that female, but not male, *16p11.2 del/+* mice have enhanced excitatory synaptic drive onto CeA neurons.

We recently identified a pathway from the CeA to the globus pallidus externa (GPe) (82), which originates predominantly from somatostatin-expressing CeA neurons that play an essential role in fear learning and anxiety (75,83,84). This pathway conveys information of the US and is critical for learning in fear conditioning (82). Importantly, optogenetic activation of the CeA-GPe pathway increases fear generalization whereby animals increase their freezing to CS-. Therefore, we sought to determine whether the GPe-projecting CeA neurons are affected by the *16p11.2* microdeletion. To this end, we used a retrograde labeling strategy whereby fluorescently conjugated CTB (cholera toxin subunit B) was injected in the GPe to label the GPe-projecting CeA neurons (Figure 6A) (see Supplemental Methods). Three days after the CTB injection, we recorded mEPSCs selectively from the CTB-labeled GPe-projecting CeA neurons in acute brain slices prepared from female or male *16p11.2 del/+* mice and their respective wild-type littermates (Figure 6A, B). Again, we found a significant interaction between sex and genotype whereby females with *16p11.2* microdeletion exhibited increased mEPSC frequency compared with wild-type littermates (Figure 6D, E). Thus, our results indicate that the *16p11.2* microdeletion caused a female-specific enhancement of

Sex-Specific Phenotypes in 16p11.2 Microdeletion Mice




Figure 2. Female *16p11.2 del/+* mice show enhanced reactivity to foot shock. **(A)** Distance traveled during 2-second shock presentations for female mice ($F_{1,50} = 14.94, p = .0003; ***p < .001$; two-way ANOVA; *16p11.2 del/+*, $n = 4$, WT, $n = 8$). **(B)** Movement velocity during 2-second shock presentations for female mice ($F_{1,50} = 2.596, p = .1135$; two-way ANOVA). **(C)** Distance traveled during 2-second shock presentations for male mice ($F_{1,50} = 1.410, p = .2407$; two-way ANOVA; *16p11.2 del/+*, $n = 7$, WT, $n = 5$). **(D)** Movement velocity during 2-second shock presentations for male mice ($F_{1,50} = 0.1467, p = .7033$; two-way ANOVA). Data are presented as mean \pm SEM. ANOVA, analysis of variance; WT, wild-type.

excitatory synaptic drive onto CeA neurons and moreover suggest dysfunction in the somatostatin-expressing CeA neurons in the CeA-GPe pathway as a potential mechanism for the increased stress susceptibility and fear generalization identified in female *16p11.2 del/+* mice.

DISCUSSION

Our results indicate that female, but not male, *16p11.2 del/+* mice have increased susceptibility to anxiety-like phenotypes following stressful life events, revealing a previously underappreciated sex-specific effect in the modulation of behavior by 16p11.2 microdeletion. Furthermore, we identify that CeA dysfunction, in particular CeA dysfunction in the CeA-GPe circuit, may underlie the female-specific behavioral phenotypes caused by the 16p11.2 microdeletion. Notably, the CeA-GPe pathway originates predominantly from somatostatin-expressing CeA neurons (82), which have been shown to have an essential role in fear learning and anxiety (75,83–86). Our results thus pinpoint a precise cellular and circuit substrate as a potential target of future therapeutics for anxiety symptoms.

These findings are consistent with the vast literature that females with ASD show distinct behavioral symptoms compared with males (22,26–29), in particular, more internalizing symptoms and emotional disturbances (30–34). Our findings are also consistent with the notion that in the general population, females have an increased prevalence of stress-related disorders, such as anxiety, depression, and posttraumatic stress disorder (40–43). Our study thus urges a careful examination of anxiety and other emotional symptoms as well as functional changes in the amygdala-

basal ganglia circuits in 16p11.2 microdeletion carriers, in particular in female carriers. In general, our study also urges sex-specific diagnostic and treatment strategies for ASD.

Three lines of evidence suggest that heightened alertness or an increase in sensitivity to aversive stimuli, or to the stimuli signaling potential threat, may underlie the increased susceptibility to anxiety-like phenotypes in female *16p11.2 del/+* mice following stressful experiences. First, *16p11.2 del/+* mice, especially females, show increased freezing when they are exposed to an unfamiliar sound, which is a sign of uncertainty or potential danger. Second, female *16p11.2 del/+* mice have enhanced reactivity to foot shocks. Third, CeA neurons in female *16p11.2 del/+* mice have enhanced sensitivity to excitatory inputs. This enhanced sensitivity may lead to heightened alertness or attention, as the CeA has been implicated in selective processing of salient information (87,88).

The CeA has central roles in the generation of fear and anxiety states (75–77,83,84,89–102). In parallel, amygdala dysfunction has been implicated in the pathogenesis of ASD. Abnormal developmental trajectory of the amygdala has been observed in ASD (103). Brain imaging studies indicate that the amygdala is enlarged precociously in children with autism (104,105) and that amygdala enlargement in autistic children is associated with anxiety symptoms (106). In addition, cellular changes in the amygdala have been reported in ASD (103). In a recent study (82), we found that a subpopulation of neurons in the CeA sends direct projections to the GPe, and the CeA-GPe pathway conveys US information and controls learning during fear conditioning. In the current study, we found




Figure 3. 16p11.2 del/+ mice have normal auditory perception. **(A)** Schematic of the behavioral task. **(B)** Performance levels across training for female mice ($F_{1,8} = 0.005112$, $p = .9448$; two-way ANOVA; 16p11.2 del/+, $n = 7$, WT, $n = 3$). **(C)** Performance levels across training for male mice ($F_{1,14} = 2.557$, $p = .1321$; two-way ANOVA; 16p11.2 del/+, $n = 9$, WT, $n = 7$). **(D)** Psychometric response curve for frequencies between 4 and 12 kHz (in log₂ values) for female mice. **(E)** Quantification of the slope of the psychometric curve (parameter p) for female mice ($p = .5878$, t test). **(F)** Quantification of the median threshold, X_0 , from the psychometric function for female mice ($p = .6465$, t test). **(G)** Average performance levels at 4 and 12 kHz for stimuli volume between 40 and 60 dB for female mice ($F_{1,8} = 0.04474$, $p = .8378$; two-way ANOVA with repeated measures). **(H)** Psychometric response curve for frequencies between 4 and 12 kHz (in log₂ values) for male mice. **(I)** Quantification of the slope of the psychometric curve (parameter p) for male mice ($p = .1713$, t test). **(J)** Quantification of the median threshold, X_0 , from the psychometric function for male mice ($p = .8607$, t test). **(K)** Average performance levels at 4 and 12 kHz for stimuli volume between 40 and 60 dB for male mice ($F_{1,14} = 0.0173$, $p = .8972$; two-way ANOVA with repeated measures). All data are presented as mean \pm SEM. ANOVA, analysis of variance; WT, wild-type.




Figure 4. Female *16p11.2 del/+* mice exhibit enhanced stress-induced anxiety-like behavior. **(A)** Time spent in the open arms of the EPM for female mice 24 hours after stress exposure ($p = .0266$; $*p < .05$; Mann-Whitney t test; female *16p11.2 del/+*, $n = 10$, female WT, $n = 8$). **(B)** Distance traveled in the EPM for female mice 24 hours after stress exposure ($p = .4082$; Mann-Whitney t test). Same mice as in panel **(A)** were used. **(C)** Movement velocity in the EPM for female mice 24 hours after stress exposure ($p = .4082$; Mann-Whitney t test). Same mice as in panel **(A)** were used. **(D)** Time spent in the open arms of the EPM for male mice 24 hours after stress exposure ($p = .3154$; Mann-Whitney t test; male *16p11.2 del/+*, $n = 8$, male WT, $n = 10$). **(E)** Distance traveled in the EPM for male mice 24 hours after stress exposure ($p = .0343$; $*p < .05$; Mann-Whitney t test). Same mice as in **(D)** were used. **(F)** Movement velocity in the EPM for male mice 24 hours after stress exposure ($p = .0266$; $*p < .05$; Mann-Whitney t test). Same mice as in **(D)** were used. **(G)** Time spent in the open arms of the EPM for female naïve mice ($p = .5629$; Mann-Whitney t test; female *16p11.2 del/+*, $n = 10$, female WT, $n = 13$). **(H)** Distance traveled in the EPM for female naïve mice ($p = .7844$; Mann-Whitney t test). Same mice as in panel **(G)** were used. **(I)** Movement velocity in the EPM for female naïve mice ($p = .1306$; Mann-Whitney t test). Same mice as in panel **(G)** were used. **(J)** Time spent in the open arms of the EPM for male naïve mice ($p = .6522$; Mann-Whitney t test; male *16p11.2 del/+*, $n = 11$, male WT, $n = 11$). **(K)** Distance traveled in the EPM for male naïve mice ($p = .6522$; Mann-Whitney t test). Same mice as in panel **(J)** are used. **(L)** Movement velocity in the EPM for male naïve mice ($p = .6522$; Mann-Whitney t test). Same mice as in panel **(J)** are used. Data are presented as mean \pm SEM. EPM, elevated plus maze; WT, wild-type.




Figure 5. Female *16p11.2 del/+* mice have increased excitatory synaptic transmission onto CeA neurons. **(A)** Schematic of the experimental design. **(B, C)** Representative mEPSC traces from CeA neurons recorded from male and female *16p11.2 del/+* **(B)** and WT **(C)** mice. **(D)** Quantification of mEPSC frequency for CeA neurons in female mice ($p = .0029$; $**p < .01$; Mann-Whitney t test; female *16p11.2 del/+*, $n = 35$ cells from 4 mice, female WT, $n = 28$ cells from 3 mice). **(E)** Quantification of mEPSC amplitude for CeA neurons in female mice ($p = .7262$; Mann-Whitney t test). Data are from the same cells as in panel **(D)**. **(F)** Quantification of mEPSC frequency for CeA neurons in male mice ($p = .2596$; Mann-Whitney t test; male *16p11.2 del/+*, $n = 21$ cells from 4 mice, male WT, $n = 14$ cells from 3 mice). **(G)** Quantification of mEPSC amplitude for CeA neurons in male mice ($p > .9999$; Mann-Whitney t test). Data are from the same cells as in panel **(F)**. Data are presented as mean \pm SEM. BA, basolateral amygdala; CeA, central amygdala; CeL, lateral division of the CeA; CeM, medial division of the CeA; LA, lateral division of the BA; mEPSC, miniature excitatory postsynaptic current; WT, wild-type.

that an enhanced excitatory drive onto GPe-projecting CeA neurons parallels the anxiety phenotypes of female *16p11.2 del/+* mice. Of note, the electrophysiology experiments in the CeA were performed on mice that were not subjected to the behavioral procedures. In addition, these mice were younger than the mice used for the behavioral experiments (see [Supplemental Methods](#)). However, because the 16p11.2 microdeletion was constitutively present from birth in these mice, it likely affected

the development of the CeA, leading to functional changes in CeA circuits even at earlier stages. We propose that early developmental and functional changes caused by the 16p11.2 microdeletion, such as those that we report here, predispose the affected individuals to anxiety-related symptoms. Our findings together strongly suggest a role of CeA-GPe circuit dysfunction in susceptibility to anxiety after stress and warrant future studies to elucidate how this circuit is involved in 16p11.2 microdeletion syndrome.




Figure 6. Female *16p11.2 del/+* mice have increased excitatory synaptic transmission onto GPe-projecting CeA neurons. **(A)** Schematic of the experimental design. CTB-555 was used to retrogradely label GPe-projecting CeA neurons. **(B, C)** Representative mEPSC traces from GPe-projecting CeA neurons recorded from male and female *16p11.2 del/+* **(B)** and WT **(C)** mice. **(D)** Quantification of mEPSC frequency for GPe-projecting CeA neurons in female mice ($p = .0482$; $*p < .05$; Mann-Whitney t test; female *16p11.2 del/+*, $n = 14$ cells from 5 mice, female WT, $n = 13$ cells from 7 mice). **(E)** Quantification of mEPSC amplitude for GPe-projecting CeA neurons in female mice ($p = .7564$; Mann-Whitney t test). Data are from the same cells as in panel **(D)**. **(F)** Quantification of mEPSC frequency for GPe-projecting CeA neurons in male mice ($p = .2017$; Mann-Whitney t test; male *16p11.2 del/+*, $n = 15$ cells from 3 mice, male WT, $n = 15$ cells from 5 mice). **(G)** Quantification of mEPSC amplitude for GPe-projecting CeA neurons in male mice ($p = .1770$; Mann-Whitney t test). Data are from the same cells as in panel **(F)**. Data are presented as mean \pm SEM. BA, basolateral amygdala; CeA, central amygdala; CeL, lateral division of the CeA; CeM, medial division of the CeA; CTB-555, cholera toxin subunit B conjugated to Alexa Fluor 555; GPe, globus pallidus externa; LA, lateral division of the BA; mEPSC, miniature excitatory postsynaptic current; WT, wild-type.

subunit B conjugated to Alexa Fluor 555; GPe, globus pallidus externa; LA, lateral division of the BA; mEPSC, miniature excitatory postsynaptic current; WT, wild-type.

ACKNOWLEDGMENTS AND DISCLOSURES

This work was supported by the National Institutes of Health (Grant Nos. R01MH101214 and R21MH114070 [to BL]), Simons Foundation (Grant No. 344904 [to BL]), National Alliance for Research on Schizophrenia and Depression (Grant No. 23169 [to BL] and Grant No. 21227 [to SA]), Cold Spring Harbor Laboratory and Northwell Health Affiliation (to BL), and Feil Family Neuroscience Endowment (to BL).

JG and BL conceived and designed the study. JG conducted the experiments and analyzed data. SA conducted the experiments with the auditory discrimination task and assisted with other electrophysiology experiments. KY identified the CeA-GPe projections and assisted with experiments. JG and BL wrote the paper with input from all authors.

We thank members of the Li laboratory for helpful discussions, Pavel Osten for providing *16p11.2 del/+* breeders, and Alea Mills for providing primers for genotyping the *16p11.2 del/+* mice.

The authors report no biomedical financial interests or potential conflicts of interest.

ARTICLE INFORMATION

From the School of Biological Sciences (JG, BL), Cold Spring Harbor Laboratory (JG, SA, KY, BL), Cold Spring Harbor, New York.

JG is currently affiliated with the Department of Psychology, University of California Los Angeles, Los Angeles, California.

SA is currently affiliated with the Laboratory of Behavioral Genetics, Brain Mind Institute, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland.

Address correspondence to Bo Li, Ph.D., at bli@cshl.edu.

Received Dec 11, 2020; revised Jan 8, 2021; accepted Jan 26, 2021.

Supplementary material cited in this article is available online at <https://doi.org/10.1016/j.bpsgos.2021.01.001>.

REFERENCES

- American Psychiatric Association (2013): Diagnostic and Statistical Manual of Mental Disorders, 5th ed. Arlington, VA: American Psychiatric Association.
- Howlin P (2000): Autism and intellectual disability: Diagnostic and treatment issues. *J R Soc Med* 93:351–355.
- Schwartz CE, Neri G (2012): Autism and intellectual disability: Two sides of the same coin. *Am J Med Genet C Semin Med Genet* 160C:89–90.
- Tonnson BL, Boan AD, Bradley CC, Charles J, Cohen A, Carpenter LA (2016): Prevalence of autism spectrum disorders among children with intellectual disability. *Am J Intellect Dev Disabil* 121:487–500.
- Antshel KM, Russo N (2019): Autism spectrum disorders and ADHD: Overlapping phenomenology, diagnostic issues, and treatment considerations. *Curr Psychiatry Rep* 21:34.
- Antshel KM, Zhang-James Y, Faraone SV (2014): The comorbidity of ADHD and autism spectrum disorder. *Expert Rev Neurother* 13:1117–1128.
- Antshel KM, Zhang-James Y, Wagner KE, Ledesma A, Faraone SV (2016): An update on the comorbidity of ADHD and ASD: A focus on clinical management. *Expert Rev Neurother* 16:1–15.
- Jang J, Matson JL, Williams LW, Tureck K, Goldin RL, Cervantes PE (2013): Rates of comorbid symptoms in children with ASD, ADHD, and comorbid ASD and ADHD. *Res Dev Disabil* 34:2369–2378.
- Leyfer OT, Folstein SE, Bacalman S, Davis NO, Dinh E, Morgan J, et al. (2006): Comorbid psychiatric disorders in children with autism: Interview development and rates of disorders. *J Autism Dev Disord* 36:849–861.
- Postorino V, Kerns CM, Vivanti G, Bradshaw J, Siracusano M, Mazzone L (2017): Anxiety disorders and obsessive-compulsive disorder in individuals with autism spectrum disorder. *Curr Psychiatry Rep* 19:92.
- Brookman-Frazee L, Stadnick N, Chlebowski C, Baker-Ericzén M, Ganger W (2018): Characterizing psychiatric comorbidity in children with autism spectrum disorder receiving publicly funded mental health services. *Autism* 22:938–952.
- White SW, Oswald D, Ollendick T, Scahill L (2009): Anxiety in children and adolescents with autism spectrum disorders. *Clin Psychol Rev* 29:216–229.
- Gotham K, Bishop SL, Hus V, Huerta M, Lund S, Buja A, et al. (2013): Exploring the relationship between anxiety and insistence on sameness in autism spectrum disorders. *Autism Res* 6:33–41.
- Andersen PN, Skogli EW, Hovik KT, Egeland J, Øie M (2015): Associations among symptoms of autism, symptoms of depression and executive functions in children with high-functioning autism: A 2 year follow-up study. *J Autism Dev Disord* 45:2497–2507.
- Davidsson M, Hult N, Gillberg C, Särneö C, Gillberg C, Billstedt E (2017): Anxiety and depression in adolescents with ADHD and autism spectrum disorders; correlation between parent- and self-reports and with attention and adaptive functioning. *Nord J Psychiatry* 71:614–620.
- Matson JL, Cervantes PE (2014): Commonly studied comorbid psychopathologies among persons with autism spectrum disorder. *Res Dev Disabil* 35:952–962.
- Hirvikoski T, Boman M, Chen Q, D’Onofrio BM, Mittendorfer-Rutz E, Lichtenstein P, et al. (2020): Individual risk and familial liability for suicide attempt and suicide in autism: a population-based study. *Psychol Med* 50:1463–1474.
- Hirvikoski T, Mittendorfer-Rutz E, Boman M, Larsson H, Lichtenstein P, Bölte S (2016): Premature mortality in autism spectrum disorder. *Br J Psychiatry* 208:232–238.
- Kirby AV, Bakian AV, Zhang Y, Bilder DA, Keeshin BR, Coon H (2019): A 20-year study of suicide death in a statewide autism population. *Autism Res* 12:658–666.
- Fombonne E (2002): Epidemiological trends in rates of autism. *Mol Psychiatry* 7(suppl 2):S4–S6.
- Baron-Cohen S (2009): Autism: The empathizing-systemizing (E-S) theory. *Ann N Y Acad Sci* 1156:68–80.
- Beggiato A, Peyre H, Maruani A, Scheid I, Rastam M, Amsellem F, et al. (2017): Gender differences in autism spectrum disorders: Divergence among specific core symptoms. *Autism Res* 10:680–689.
- Knickmeyer RC, Wheelwright S, Baron-Cohen SB (2008): Sex-typical play: Masculinization/defeminization in girls with an autism spectrum condition. *J Autism Dev Disord* 38:1028–1035.
- Kopp S, Beckung E, Gillberg C (2010): Developmental coordination disorder and other motor control problems in girls with autism spectrum disorder and/or attention-deficit/hyperactivity disorder. *Res Dev Disabil* 31:350–361.
- Szatmari P, Liu X, Goldberg J, Zwaigenbaum L, Paterson AD, Woodbury-Smith M, et al. (2012): Sex differences in repetitive stereotyped behaviors in autism: Implications for genetic liability. *Am J Med Genet B Neuropsychiatr Genet* 159B:5–12.
- Dean M, Harwood R, Kasari C (2017): The art of camouflage: Gender differences in the social behaviors of girls and boys with autism spectrum disorder. *Autism* 21:678–689.
- Head AM, McGillivray JA, Stokes MA (2014): Gender differences in emotionality and sociability in children with autism spectrum disorders. *Mol Autism* 5:19.
- Hiller RM, Young RL, Weber N (2014): Sex differences in autism spectrum disorder based on DSM-5 criteria: Evidence from clinician and teacher reporting. *J Abnorm Child Psychol* 42:1381–1393.
- Werling DM, Geschwind DH (2013): Sex differences in autism spectrum disorders. *Curr Opin Neurol* 26:146–153.
- Horiuchi F, Oka Y, Uno H, Kawabe K, Okada F, Saito I, et al. (2014): Age- and sex-related emotional and behavioral problems in children with autism spectrum disorders: Comparison with control children. *Psychiatry Clin Neurosci* 68:542–550.
- Kreiser NL, White SW (2014): ASD in females: Are we overstating the gender difference in diagnosis? *Clin Child Fam Psychol Rev* 17:67–84.
- Rynkiewicz A, Łucka I (2018): Autism spectrum disorder (ASD) in girls. Co-occurring psychopathology. Sex differences in clinical manifestation. *Psychiatr Pol* 52:629–639.
- Rynkiewicz A, Schuller B, Marchi E, Piana S, Camurri A, Lassalle A, Baron-Cohen S (2016): An investigation of the ‘female camouflage

- effect' in autism using a computerized ADOS-2 and a test of sex/gender differences. *Mol Autism* 7:10.
34. Solomon M, Miller M, Taylor SL, Hinshaw SP, Carter CS (2012): Autism symptoms and internalizing psychopathology in girls and boys with autism spectrum disorders. *J Autism Dev Disord* 42:48–59.
 35. Kalyva E (2009): Comparison of eating attitudes between adolescent girls with and without Asperger syndrome: Daughters' and mothers' reports. *J Autism Dev Disord* 39:480–486.
 36. Lai MC, Lombardo MV, Baron-Cohen S (2014): Autism. *Lancet* 383:896–910.
 37. Hartley SL, Sikora DM (2009): Sex differences in autism spectrum disorder: An examination of developmental functioning, autistic symptoms, and coexisting behavior problems in toddlers. *J Autism Dev Disord* 39:1715.
 38. Giarelli E, Wiggins LD, Rice CE, Levy SE, Kirby RS, Pinto-Martin J, Mandell D (2010): Sex differences in the evaluation and diagnosis of autism spectrum disorders among children. *Disabil Health J* 3:107–116.
 39. Bölte S, Duketis E, Poustka F, Holtmann M (2011): Sex differences in cognitive domains and their clinical correlates in higher-functioning autism spectrum disorders. *Autism* 15:497–511.
 40. Olf M (2017): Sex and gender differences in post-traumatic stress disorder: An update. *Eur J Psychotraumatol* 8(suppl 4):1351204.
 41. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995): Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 52:1048–1060.
 42. Breslau N (2002): Gender differences in trauma and posttraumatic stress disorder. *J Genet Specif Med* 5:34–40.
 43. Tolin DF, Foa EB (2006): Sex differences in trauma and posttraumatic stress disorder: A quantitative review of 25 years of research. *Psychol Bull* 132:959–992.
 44. Hughes RN (2007): Sex does matter: Comments on the prevalence of male-only investigations of drug effects on rodent behaviour. *Behav Pharmacol* 18:583–589.
 45. Beery AK, Zucker I (2011): Sex bias in neuroscience and biomedical research. *Neurosci Biobehav Rev* 35:565–572.
 46. Marshall CR, Noor A, Vincent JB, Lionel AC, Feuk L, Skaug J, *et al.* (2008): Structural variation of chromosomes in autism spectrum disorder. *Am J Hum Genet* 82:477–488.
 47. Weiss LA, Shen Y, Korn JM, Arking DE, Miller DT, Fossdal R, *et al.* (2008): Association between microdeletion and microduplication at 16p11.2 and autism. *New Engl J Med* 358:667–675.
 48. Kumar RA, KaraMohamed S, Sudi J, Conrad DF, Brune C, Badner JA, *et al.* (2007): Recurrent 16p11.2 microdeletions in autism. *Hum Mol Genet* 17:628–638.
 49. Sanders SJ, Ercan-Sencicek AG, Hus V, Luo R, Murtha MT, Moreno-De-Luca D, *et al.* (2011): Multiple recurrent de novo CNVs, including duplications of the 7q11.23 Williams syndrome region, are strongly associated with autism. *Neuron* 70:863–885.
 50. Chen CH, Chen HI, Chien WH, Li LH, Wu YY, Chiu YN, *et al.* (2017): High resolution analysis of rare copy number variants in patients with autism spectrum disorder from Taiwan. *Sci Rep* 7:11919.
 51. Sebat J, Lakshmi B, Malhotra D, Troge J, Lese-Martin C, Walsh T, *et al.* (2007): Strong association of de novo copy number mutations with autism. *Science* 316:445–449.
 52. Bjijsma EK, Gijsbers AC, Schuurs-Hoeijmakers JH, van Haeringen A, Franssen van de Putte DE, Anderlid BM, *et al.* (2009): Extending the phenotype of recurrent rearrangements of 16p11.2: Deletions in mentally retarded patients without autism and in normal individuals. *Eur J Med Genet* 52:77–87.
 53. Fernandez BA, Roberts W, Chung B, Weksberg R, Meyn S, Szatmari P, *et al.* (2010): Phenotypic spectrum associated with de novo and inherited deletions and duplications at 16p11.2 in individuals ascertained for diagnosis of autism spectrum disorder. *J Med Genet* 47:195.
 54. Shinawi M, Liu P, Kang SH, Shen J, Belmont JW, Scott DA, *et al.* (2010): Recurrent reciprocal 16p11.2 rearrangements associated with global developmental delay, behavioural problems, dysmorphism, epilepsy, and abnormal head size. *J Med Genet* 47:332–341.
 55. Al-Kateb H, Khanna G, Filges I, Hauser N, Grange DK, Shen J, *et al.* (2014): Scoliosis and vertebral anomalies: Additional abnormal phenotypes associated with chromosome 16p11.2 rearrangement. *Am J Med Genet A* 164:1118–1126.
 56. Steinman KJ, Spence SJ, Ramocki MB, Proud MB, Kessler SK, Marco EJ, *et al.* (2016): 16p11.2 deletion and duplication: Characterizing neurologic phenotypes in a large clinically ascertained cohort. *Am J Med Genet A* 170:2943–2955.
 57. Zufferey F, Sherr EH, Beckmann ND, Hanson E, Maillard AM, Hippolyte L, *et al.* (2012): A 600 kb deletion syndrome at 16p11.2 leads to energy imbalance and neuropsychiatric disorders. *J Med Genet* 49:660–668.
 58. Horev G, Ellegood J, Lerch JP, Son YE, Muthuswamy L, Vogel H, *et al.* (2011): Dosage-dependent phenotypes in models of 16p11.2 lesions found in autism. *Proc Natl Acad Sci U S A* 108:17076–17081.
 59. Portmann T, Yang M, Mao R, Panagiotakos G, Ellegood J, Dolen G, *et al.* (2014): Behavioral abnormalities and circuit defects in the basal ganglia of a mouse model of 16p11.2 deletion syndrome. *Cell Rep* 7:1077–1092.
 60. Arbogast T, Ouagazzal AM, Chevalier C, Kopanitsa M, Afinowi N, Migliavacca E, *et al.* (2016): Reciprocal effects on neurocognitive and metabolic phenotypes in mouse models of 16p11.2 deletion and duplication syndromes. *PLoS Genet* 12:e1005709.
 61. Yang M, Lewis FC, Sarvi MS, Foley GM, Crawley JN (2015): 16p11.2 deletion mice display cognitive deficits in touchscreen learning and novelty recognition tasks. *Learn Mem* 22:622–632.
 62. Yang M, Mahrt EJ, Lewis F, Foley G, Portmann T, Dolmetsch RE, *et al.* (2015): 16p11.2 deletion syndrome mice display sensory and ultrasonic vocalization deficits during social interactions. *Autism Res* 8:507–521.
 63. Angelakos CC, Watson AJ, O'Brien WT, Krainock KS, Nickl-Jockschat T, Abel T (2017): Hyperactivity and male-specific sleep deficits in the 16p11.2 deletion mouse model of autism. *Autism Res* 10:572–584.
 64. Grissom NM, McKee SE, Schoch H, Bowman N, Havekes R, O'Brien WT, *et al.* (2017): Male-specific deficits in natural reward learning in a mouse model of neurodevelopmental disorders. *Mol Psychiatry* 23:544–555.
 65. Walsh JJ, Christoffel DJ, Heifets BD, Ben-Dor GA, Selimbeyoglu A, Hung LW, *et al.* (2018): 5-HT release in nucleus accumbens rescues social deficits in mouse autism model. *Nature* 560:589–594.
 66. Rein B, Yan Z (2020): 16p11.2 copy number variations and neurodevelopmental disorders. *Trends Neurosci* 43:886–901.
 67. Lynch JF 3rd, Ferri SL, Angelakos C, Schoch H, Nickl-Jockschat T, Gonzalez A, *et al.* (2020): Comprehensive behavioral phenotyping of a 16p11.2 Del mouse model for neurodevelopmental disorders. *Autism Res* 13:1670–1684.
 68. Pucilowska J, Vithayathil J, Tavares EJ, Kelly C, Karlo JC, Landreth GE (2015): The 16p11.2 deletion mouse model of autism exhibits altered cortical progenitor proliferation and brain cytoarchitecture linked to the ERK MAPK pathway. *J Neurosci* 35:3190–3200.
 69. Tian D, Stoppel LJ, Heynen AJ, Lindemann L, Jaeschke G, Mills AA, Bear MF (2015): Contribution of mGluR5 to pathophysiology in a mouse model of human chromosome 16p11.2 microdeletion. *Nat Neurosci* 18:182–184.
 70. Hudac CM, Bove J, Barber S, Duyzend M, Wallace A, Martin CL, *et al.* (2020): Evaluating heterogeneity in ASD symptomatology, cognitive ability, and adaptive functioning among 16p11.2 CNV carriers. *Autism Res* 13:1300–1310.
 71. Panzini CM, Ehlinger DG, Alchahin AM, Guo Y, Commons KG (2017): 16p11.2 deletion syndrome mice persevere with active coping response to acute stress—rescue by blocking 5-HT2A receptors. *J Neurochem* 143:708–721.
 72. Dalla C, Shors TJ (2009): Sex differences in learning processes of classical and operant conditioning. *Physiol Behav* 97:229–238.
 73. Gruene TM, Flick K, Stefano A, Shea SD, Shansky RM (2015): Sexually divergent expression of active and passive conditioned fear responses in rats. *Elife* 4:e11352.

Sex-Specific Phenotypes in 16p11.2 Microdeletion Mice

74. Dunsmoor JE, Paz R (2015): Fear generalization and anxiety: Behavioral and neural mechanisms. *Biol Psychiatry* 78:336–343.
75. Ahrens S, Wu MV, Furlan A, Hwang GR, Paik R, Li H, *et al.* (2018): A central extended amygdala circuit that modulates anxiety. *J Neurosci* 38:5567–5583.
76. Shackman AJ, Fox AS (2016): Contributions of the central extended amygdala to fear and anxiety. *J Neurosci* 36:8050–8063.
77. Fox AS, Oler JA, Shelton SE, Nanda SA, Davidson RJ, Roseboom PH, Kalin NH (2012): Central amygdala nucleus (Ce) gene expression linked to increased trait-like Ce metabolism and anxious temperament in young primates. *Proc Natl Acad Sci U S A* 109:18108–18113.
78. Ahrens S, Jaramillo S, Yu K, Ghosh S, Hwang GR, Paik R, *et al.* (2015): ErbB4 regulation of a thalamic reticular nucleus circuit for sensory selection. *Nat Neurosci* 18:104–111.
79. Varodayan FP, Khom S, Patel RR, Steinman MQ, Hedges DM, Oleata CS, *et al.* (2018): Role of TLR4 in the modulation of central amygdala GABA transmission by CRF following restraint stress. *Alcohol Alcohol* 53:642–649.
80. Varodayan FP, Minnig MA, Steinman MS, Oleata CS, Riley MW, Sabino V, Roberto M (2020): PACAP regulation of central amygdala GABAergic synapses is altered by restraint stress. *Neuropharmacology* 168:107752.
81. Zimprich A, Garrett L, Deussing JM, Wotjak CT, Fuchs H, Gailus-Durner V, *et al.* (2014): A robust and reliable non-invasive test for stress responsivity in mice. *Front Behav Neurosci* 8:125.
82. Giovanniello J, Yu K, Furlan A, Nachtrab GT, Sharma R, Chen X, Li B (2020): A central amygdala-globus pallidus circuit conveys unconditioned stimulus-related information and controls fear learning. *J Neurosci* 40:9043–9054.
83. Li H, Penzo MA, Taniguchi H, Kopec CD, Huang ZJ, Li B (2013): Experience-dependent modification of a central amygdala fear circuit. *Nat Neurosci* 16:332.
84. Penzo MA, Robert V, Tucciarone J, Bundel DD, Wang M, Aelst LV, *et al.* (2015): The paraventricular thalamus controls a central amygdala fear circuit. *Nature* 519:455.
85. Shrestha P, Shan Z, Mamcarz M, Ruiz KSA, Zerihoun AT, Juan CY, *et al.* (2020): Amygdala inhibitory neurons as loci for translation in emotional memories. *Nature* 586:407–411.
86. Hartley ND, Gaulden AD, Báldi R, Winters ND, Salimando GJ, Rosas-Vidal LE, *et al.* (2019): Dynamic remodeling of a basolateral-to-central amygdala glutamatergic circuit across fear states. *Nat Neurosci* 22:2000–2012.
87. Calu DJ, Roesch MR, Haney RZ, Holland PC, Schoenbaum G (2010): Neural correlates of variations in event processing during learning in central nucleus of amygdala. *Neuron* 68:991–1001.
88. Roesch MR, Esber GR, Li J, Daw ND, Schoenbaum G (2012): Surprise! Neural correlates of Pearce-Hall and Rescorla-Wagner coexist within the brain. *Eur J Neurosci* 35:1190–1200.
89. Etkin A, Wager TD (2007): Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 164:1476–1488.
90. Wager DT, Barrett FL, Bliss-Moreau E, Lindquist KA, Duncan S, Kober H, *et al.* (2008): The Neuroimaging of Emotion. In: Lewis M, Haviland-Jones JM, Barrett LF, editors. *Handbook of Emotions*, 3rd ed. New York: The Guilford Press, 249–271.
91. Walker DL, Davis M (2008): Role of the extended amygdala in short-duration versus sustained fear: A tribute to Dr. Lennart Heimer. *Brain Struct Funct* 213:29–42.
92. Davis M, Walker DL, Miles L, Grillon C (2010): Phasic vs sustained fear in rats and humans: Role of the extended amygdala in fear vs anxiety. *Neuropsychopharmacology* 35:105–135.
93. Mobbs D, Yu R, Rowe JB, Eich H, FeldmanHall O, Dalgleish T (2010): Neural activity associated with monitoring the oscillating threat value of a tarantula. *Proc Natl Acad Sci U S A* 107:20582–20586.
94. Jennings JH, Sparta DR, Stamatakis AM, Ung RL, Pleil KE, Kash TL, Stuber GD (2013): Distinct extended amygdala circuits for divergent motivational states. *Nature* 496:224–228.
95. Kim SY, Adhikari A, Lee SY, Marshel JH, Kim CK, Mallory CS, *et al.* (2013): Diverging neural pathways assemble a behavioural state from separable features in anxiety. *Nature* 496:219–223.
96. Andreatta M, Glotzbach-Schoon E, Mühlberger A, Schulz SM, Wiemer J, Pauli P (2015): Initial and sustained brain responses to contextual conditioned anxiety in humans. *Cortex* 63:352–363.
97. Calhoun GG, Tye KM (2015): Resolving the neural circuits of anxiety. *Nat Neurosci* 18:1394–1404.
98. Fox AS, Oler JA, Tromp DPM, Fudge JL, Kalin NH (2015): Extending the amygdala in theories of threat processing. *Trends Neurosci* 38:319–329.
99. Tovote P, Fadok JP, Lüthi A (2015): Neuronal circuits for fear and anxiety. *Nat Rev Neurosci* 16:317–331.
100. Gungor NZ, Paré D (2016): Functional heterogeneity in the bed nucleus of the stria terminalis. *J Neurosci* 36:8038–8049.
101. Marcinkiewicz CA, Mazonne CM, D'Agostino G, Halladay LR, Hardaway JA, DiBerto JF, *et al.* (2016): Serotonin engages an anxiety and fear-promoting circuit in the extended amygdala. *Nature* 537:97–101.
102. Yu K, Ahrens S, Zhang X, Schiff H, Ramakrishnan C, Fenno L, *et al.* (2017): The central amygdala controls learning in the lateral amygdala. *Nat Neurosci* 20:1680–1685.
103. Amaral DG, Schumann CM, Nordahl CW (2008): Neuroanatomy of autism. *Trends Neurosci* 31:137–145.
104. Sparks BF, Friedman SD, Shaw DW, Aylward EH, Echelard D, Artru AA, *et al.* (2002): Brain structural abnormalities in young children with autism spectrum disorder. *Neurology* 59:184–192.
105. Schumann CM, Hamstra J, Goodlin-Jones BL, Lotspeich LJ, Kwon H, Buonocore MH, *et al.* (2004): The amygdala is enlarged in children but not adolescents with autism; the hippocampus is enlarged at all ages. *J Neurosci* 24:6392–6401.
106. Juranek J, Filipek PA, Berenji GR, Modahl C, Osann K, Spence MA (2006): Association between amygdala volume and anxiety level: Magnetic resonance imaging (MRI) study in autistic children. *J Child Neurol* 21:1051–1058.